

TRYPANOSOMA AVIUM INCIDENCE, PATHOGENICITY AND RESPONSE TO MELARSOMINE IN FALCONS FROM KUWAIT

W. TARELLO*

Summary:

Epidemiological and clinical studies on *Trypanosoma avium* are lacking in the Middle East. The aims of this study were to determine the *T. avium* incidence in falcons from Kuwait, report clinical signs and find an effective therapy. Blood smears from 921 diseased and 56 healthy falcons were examined between May 2003 and April 2004. 12 birds (1.3 %) were found infected by *T. avium* and ten of these were treated with melarsomine (Cymelarsan®) at a dosage of 0.25 mg/kg intramuscularly for four days. All affected birds presented clinical signs, including incapacity of flying high, poor appetite, lethargy, loosing weight, weakness, dyspnoea and death. Signs disappeared within 1-7 days after administration of melarsomine. Trypomastigotes were not detected in blood smears made 1-7 days after the end of therapy. This study suggests that *T. avium* induces disease in falcons and that melarsomine can be an effective therapy eliminating both clinical signs and circulating trypomastigotes.

KEY WORDS : *Trypanosoma avium*, falcon, bird of prey, pathogenicity, chemotherapy, melarsomine, Kuwait.

Résumé : INCIDENCE DE *TRYPANOSOMA AVIUM*, SYMPTOMATOLOGIE ET RÉPONSE AU TRAITEMENT AVEC LA MÉLARSOMINE CHEZ LES FAUCONS AU KOWEIT

Au Moyen-Orient, il y a un manque d'études épidémiologiques et cliniques sur *Trypanosoma avium*. La présente étude avait pour objectifs de déterminer l'incidence de *T. avium* chez les faucons au Koweït, de décrire les signes cliniques et de trouver une thérapie efficace. Du mois de mai 2003 au mois d'avril 2004, les frottis sanguins de 921 faucons malades et de 56 faucons sains ont été examinés. 12 oiseaux (1,3 %) étaient porteurs de *T. avium* et dix ont été soumis à un traitement intramusculaire de mélarsomine (Cymelarsan®) à la posologie de 0,25 mg/kg par jour pendant quatre jours. Tous les oiseaux atteints présentaient des signes cliniques incluant une incapacité au haut vol, un manque d'appétit, une léthargie, une perte de poids, un épuisement général, une difficulté respiratoire et, pour certains, la mort. Les signes cliniques ont disparu en un à sept jours et on n'a plus trouvé de trypomastigotes dans le sang une semaine après la thérapie. Cette étude suggère que *T. avium* provoque une maladie chez les faucons de chasse et que la mélarsomine peut être un traitement de choix en éliminant les trypomastigotes et les signes cliniques.

MOTS CLÉS : *Trypanosoma avium*, faucon, oiseau de proie, symptomatologie, chimiothérapie, mélarsomine, Koweït.

Trypanosoma avium is a worldwide distributed flagellate parasite of birds (Peirce, 2003) transmitted by mosquitoes and ornithophilic flies (Votycka *et al.*, 2002; Votycka & Svobodova, 2004) showing poor host specificity and wide geographical distribution (Sehgal *et al.*, 2001). These organisms are highly pleomorphic and intense parasitemias are not uncommon (Peirce, 2003). There are very few data available on the prevalence of *Trypanosoma avium* in Falconiformes (Kirkpatrick & Lauer, 1985; Munoz *et al.*, 1999) and a specific study has never been carried out in the Middle East. Current scientific knowledge suggests that no pathogenicity is associated with avian trypanosomosis (Peirce, 2003). However, preliminary

observations made during the initial part of this study indicate that clinical signs are present in captive falcons infected by *T. avium*. This paper reports the incidence of *T. avium* in a large population of sick falcons, the clinical signs observed, and the response to melarsomine (Cymelarsan®, Merial), a trypanocide drug that has never previously been used in birds infected by *T. avium*.

MATERIALS AND METHODS

From early May 2003 to the end of April 2004, examinations for the presence of *Trypanosoma avium* were made on Diff-Quick-stained blood smears from 921 diseased birds of prey and 56 healthy control falcons, belonging to three species (*Falco peregrinus*, *Falco cherrug* and *Falco rusticolus*).

Blood samples were collected from the brachial vein. All falcons were used for hunting, a category given

* International Veterinary Hospital, P.O. Box 9275, 61003 Ahmadi, Kuwait.
Tel.: 00 965 3261421 – Fax: 00 965 3261392
E-mail: wtarello@yahoo.it

constant special care in the Middle East. Consequently, reports on their health status and clinical signs were precise. Information evaluated included signalement, origin, date of consultation, clinical signs, intensity of infection (+ = low, ++ = moderate, +++ = high, ++++ = very high) and melarsomine (Cymelarsan®, Merial) therapy outcomes (Table I). Melarsomine, a registered medication for the treatment of *Trypanosoma evansi* in camels ("surra"), was injected by IM route at the recommended daily dose of 0.25 mg/kg (Touratier, 1992) for four consecutive days. Clinical and haematological outcomes were checked within 1-7 days after the end of therapy.

RESULTS

Twelve falcons out of 921 (1,3 %) were infected (Table I). None of the 56 healthy controls was carrying *T. avium*. In total, 977 birds of prey were examined. Trypanosome infections were detected between August and January, but not between February and July. Clinical signs were present in all affected raptors, and no concurrent diseases could be detected in any of them. Sudden death occurred in one case. Signs more often reported were incapacity of flying high (7; 58.3 %), poor appetite (6; 50 %), lethargy (5; 41.7 %), loose of weight (5; 41.7 %),

weakness (3; 25 %) dyspnoea (2; 16.7 %) and a cessation of molting (2; 16.7 %). Within seven days after the end of therapy, the ten treated birds presented complete clinical recovery with the disappearance of trypanomastigotes (Table I).

DISCUSSION

Scattered past reports have pointed out that *T. avium* can cause fatal infections in birds (Baker, 1976; Mungomba *et al.*, 1989). One falcon in the study group died before starting treatment, apparently confirming those previous observations. According to the recent literature, there was no evidence to indicate that *T. avium* is pathogenic (Peirce, 2003; Votypka *et al.*, 2002).

This assumption does not seem to be based on verifiable facts but on the lack of direct research, since today only morphological (Yurchenko *et al.*, 1999) and epidemiological informations (Kirkpatrick & Lauer, 1985; Munoz *et al.*, 1999) are available. Most of previous studies were carried out on wild animals captured and released after sampling (Rintamaki *et al.*, 1999; Deviche *et al.*, 2001; Holstad *et al.*, 2003). Under such field conditions pathogenic effects are hard to assess. In domestic or semi-domesticated animals living in proximity to human beings, such as captive falcons,

No.	Species, sex, age, weight, origin and date of visit	Duration of the disease	Clinical signs and intensity of infection (from + to ++++)	Therapy outcomes
1	Saker, F, 4 years, 975 gr, Kuwait, 25 August 2003	1 month	Poor appetite, lethargy, stopped molting (++++)	Dead before starting the treatment
2	Peregrine, M, 1 year, 640 gr, Pakistan, 26 August 2003	1 week	Poor appetite, lethargy, loosing weight (++++)	Clinical and haematological recovery
3	Saker, F, 4 years, 1114 gr, Kuwait, 29 August 2003	3-5 days	Lethargy (++)	Clinical and haematological recovery
4	Saker, F, 3 years, 985 gr, Russia, 29 September 2003	2 months	Poor appetite, stopped molting, loosing weight (++)	Clinical and haematological recovery
5	Peregrine, M, 4 years, 841 gr, Kuwait, 28 October 2003	10 days	Poor appetite, dyspnoea, weakness (+++)	Clinical and haematological recovery
6	Saker, M, 2 years, 850 gr, Kuwait, 15 November 2003	1 week	Weakness, not flying high, blood in the stool (+++)	Clinical and haematological recovery
7	Peregrine, M, 1 year, 608 gr, Iraq, 26 November 2003	1 week	Lethargy, loosing weight (++++)	Clinical and haematological recovery
8	Saker, M, 4 years, 900 gr, Kuwait, 26 November 2003	10 days	Poor appetite, not flying high, loosing weight (+++)	Clinical and haematological recovery
9	Saker, F, 1 year, 1068 gr, Kuwait, 6 December 2003	1 month	Weakness, not flying high, dyspnoea (+)	Treatment was denied
10	Peregrine, M, 1 year, 926 gr, Kuwait, 10 January 2004	1week	Not flying high, haematoma on the back (++++)	Clinical and haematological recovery
11	Saker, M, 1 year, 790 gr, Iraq, 21 January 2004	1 month	Poor appetite, lethargy, not flying high (++++)	Clinical and haematological recovery
12	Saker, F, 3 years, 1,000 gr, Kuwait, 29 January 2004	1 month	Weakness, loosing weight, not flying high (++++)	Clinical and haematological recovery

Table I. – *Trypanosoma avium* and response to therapy in 12 falcons from Kuwait.

pathogenic effects are easier to detect and to describe. This seems to be the first attempt to study clinical signs and the response to therapy of *T. avium* infection in captive birds of prey. The results obtained indicate the constant presence of pathogenic effects and 9 % mortality rate (Table I). Fifty-six healthy controls proved negative for *T. avium*, leading to the exclusion of healthy carriers in the study group. The disappearance of clinical signs and of trypanosomes shortly after trypanocide therapy with melarsomine indirectly confirmed that the pathogenic signs were linked to the presence of trypanosomes.

The identification of trypomastigotes in blood smears is the diagnostic test for trypanosomiasis suggested by the OIE (OIE, 2000). Flagellate bodies observed in the blood from the 12 falcons showed shapes and sizes in accordance with the morphological features of *Trypanosoma avium* (Peirce, 2003). Although blood samples were always obtained from the brachial vein, it is acknowledged that there is no significant difference between *T. avium* prevalence in blood collected from the brachial vein or deep circulation (Holstad *et al.*, 2003). Thus, the incidence rate of 1.3 % obtained in this study should not be controversial with regard to the sampling site, although blood cultures may reveal a much higher prevalence. As an example, 1.2 % of 259 birds of prey from southern New Jersey showed circulating *Trypanosoma* spp., while blood cultures from 142 of these raptors revealed a prevalence of 41.5 % (Kirkpatrick & Lauer, 1985). It has been noted that the examination of stained blood smears will only show trypanosomes when present in the host at high levels (Peirce, 2003). This might explain the observation of clinical signs in infected birds, since it seems reasonable to assume that the higher the number of parasites the more pathogenic is their action.

The 1.3 % incidence reported here is low if compared with studies from areas at higher latitudes (Rintamaki *et al.*, 1999; Deviche *et al.*, 2001; Holstad *et al.*, 2003), taking also into account that the population examined includes 921 'diseased' falcons and not one healthy falcon.

However, results from studies carried out on free-ranging falcons admitted to wildlife rescue centres, accordingly to be considered 'diseased', are overlapping. In fact, 190 diseased falcons from Spain were found trypanosome-free (Munoz *et al.*, 1999), and only 1.2 % of 259 diseased birds of prey from USA were found to carry trypanosomes (Kirkpatrick & Lauer, 1985). Unfortunately, clinical signs were not recorded and no therapy was suggested.

The abundance of vectors enhances parasite transmission during favouring seasons (Rintamaki *et al.*, 1999). This observation is in agreement with the results of this one-year study since all cases were recorded

between August and January and none between February and July (Table I), apparently indicating a seasonal occurrence.

REFERENCES

- BAKER J.R. Biology of trypanosomes of birds. *In: Biology of the Kinetoplastida*. Lumsden W.H.R. & Evans D.A. (eds). Academic Press, New York, 1976, 131-174.
- DEVICHE P., GREINER E.C. & MANTECA X. Interspecific variability of prevalence in blood parasites of adult passerine birds during the breeding season in Alaska. *Journal of Wildlife Diseases* 2001, 37, 28-35.
- HOLSTAD P.R., ANWAR A., IEZHOVA T. & SKORPING A. Standard sampling techniques underestimate prevalence of avian hematozoa in willow ptarmigan (*Lagopus lagopus*). *Journal of Wildlife Diseases* 2003, 39, 354-358.
- KIRKPATRICK C.E. & LAUER D.M. Hematozoa of raptors from southern New Jersey and adjacent areas. *Journal of Wildlife Diseases*, 1985, 21, 1-6.
- MUNGOMBA L.M., MOLYNEAUX D.H. & WALLBANKS K.R. Host-parasite relationship of *Trypanosoma corvi* in *Ornithomyia avicularia*. *Parasitology Research*, 1989, 75, 167-174.
- MUNOZ E., FERRER D., MOLINA R. & ADLARD R.D. Prevalence of haematozoa in birds of prey in Catalonia, north-east Spain. *Veterinary Record*, 1999, 144, 632-636.
- OIE. Trypanosomiasis. *In: Manual of standard for diagnostic tests and vaccines 2000*. www.oie.int/eng/normes/manual. Accessed August 20, 2004.
- PEIRCE M.A. Haematozoa. *In: Avian medicine*. Samour J. (ed.). Elsevier Science, Edinburgh, 2003, 245-252.
- RINTAMAKI P.T., HUHTA E., JOKIMAKI J. & SQUIRES-PARSONS D. Leucocytozoonosis and trypanosomiasis in redstarts in Finland. *Journal of Wildlife Diseases*, 1999, 35, 603-607.
- SEHGAL R.N., JONES H.I. & SMITH T.B. Host specificity and incidence of *Trypanosoma* in some African rainforest birds: a molecular approach. *Molecular Ecology*, 2001, 10, 2319-2327.
- VOTYPKA J., OBORNIK M., VOLF P., SVOBODOVA M. & LUKES J. *Trypanosoma avium* of raptors (Falconiformes): phylogeny and identification of vectors. *Parasitology*, 2002, 125, 253-263.
- VOTYPKA J. & SVOBODOVA M. *Trypanosoma avium*: experimental transmission from black flies to canaries. *Parasitology Research*, 2004, 92, 147-51.
- TOURATIER L. Eleventh international meeting on *Trypanosoma evansi*: report of the Working Group. Paris, 17 May 1990. *Revue Scientifique et Technique*, 1992, 11, 275-304.
- YURCHENKO V., HOBZA R., BENADA O. & LUKES J. *Trypanosoma avium*: large minicircles in the kinetoplast DNA. *Experimental Parasitology*, 1999, 92, 215-218.

Reçu le 16 septembre 2004
 Accepté le 14 décembre 2004