

Tables S2: Nymphal infection prevalence (NIP)

	Robertsau	Pourtalès	Orangerie	Botanical	Citadelle	Schulmeister	Neudorf	Rohrschollen	Total
Infected nymphs	67	12	3	1	0	0	51	48	182
Non-infected	117	67	21	9	4	1	142	114	475
Total	184	79	24	10	4	1	193	162	657
NIP %	36.4	15.2	12.5	10.0	/	/	26.4	29.6	27.7
95% CI	29.5–43.9	8.4–25.4	3.3–33.5	0.5–45.9			20.5–33.3	22.9–37.4	24.3–31.3

Table S2a: Nymphal infection prevalence (NIP) for the 8 sites.

/: insufficient data

	North	City	South	Total
Infected nymphs	79	4	99	182
Non-infected	184	35	256	475
Total	263	39	355	657
NIP %	30.0	10.3	27.9	27.7
95% CI	24.8–35.8	4.1–23.6	23.5–32.8	24.3–31.3

Table S2b: Nymphal infection prevalence (NIP) for the 3 zones (ecosystems).

	March	April	May	June	Total
Infected nymphs	42	29	57	54	182
Non-infected	121	144	84	126	475
Total	163	173	141	180	657
NIP %	25.8	16.8	40.4	30.0	27.7
95% CI	19.4–33.3	11.7–23.4	32.3–49.0	23.5–37.3	24.3–31.3

Table S2c: Nymphal infection prevalence (NIP) for the 4 months.