mainly investigated, although other parts of the country were visited to a lesser extent. Onchocerciasis was found only in villages along the former two rivers. No infected individuals were recorded from 987 people sampled at thirty-five localities in the regions of Cacheu, Oio and Tombali outside the known foci.

Of 516 people examined in six villages along the Geba river 102 (19 %) were infected with the disease. The data show that the disease was more prevalent in the older age groups. Skin densities were relatively low with the majority of infected individuals having less than 15 mff/mg.

Investigations in villages along the Corubal river showed that 316 (37 %) of 833 individuals examined were infected. An analysis of the disease in relation to age and sex shows that although larger numbers of older people were infected the disease was also significant in younger males. Skin densities were also higher than for individuals from villages along the Geba river.

Typical symptoms of onchocerciasis were seen in infected individuals and 48 of the infected people were blind (31 males and 17 females). In most cases this was due to the presence of microfilariae in the eye.

The data presented indicate that the more important of the two foci is that of the Corubal river. Irrigation schemes along the upper reaches of the Geba river have drastically reduced water flow and transmission is believed to be only sporadic or even absent because of the paucity of *S. damnosum* s.l. in this area.

SIMULIIDAE

The distribution of Simuliidae in Guinea Bissau was investigated (Grácio *et al.*, in press) and identification of the cytospecies of the *Simulium damnosum* complex made at localities within and adjacent to the Corubal focus (Charalambous *et al.*, in press).

Only *S. sirbanum* and *S. konkoureense* were found along the Corubal river between March 1990 and November 1992. The former species predominated at localities along the upper reaches and the largely zoophilic *S. konkoureense* was the main cytoform along the lower reaches of the river. Morphometric analysis of cytotyped larvae showed these two cytota of *S. damnosum* could be separated using the number of rows of hooks in the posterior circlet and the number of hypostomial setae. Unfortunately, this differentiation of the two cytotypes contained a large percentage of overlap so that unequivocal identifications could only be made for specimens in the extremes of the combined range (Lowry *et al.*, in prep.).

Entomological surveys showed the presence of the following simuliid species associated with the Geba and Corubal river basins : *Simulium alcocki*, *S. garmsi*, *S. djallonense*, *S. cervicornutum*, *S. unicornutum*, *S. ruficornis*, *S. adersi*, *S. tridens* and *S. bargreavesi*.

BITING BEHAVIOUR AND HOST CAPACITIES

Adult *S. sirbanum* and *S. konkoureense* were collected biting man and large numbers of the former cytospecies were experimentally infected with *O. volvulus* to com-

pare filarial development in the two cytospecies. Insufficient numbers of *S. konkoureense* precluded the completion of the experiment.

REFERENCES

- ABREU M.M., MARINI A. : Contribuição para o estudo da oncocercose na Guiné Portuguesa (Foco do Rio Corubal). *Bol. Cult. Guiné Port.*, 1964, 19, 433-453 (also published in *Rev. Est. Ger. Univ. Moçambique*, 1964, 1, 3-23).
- ABREU M.M., MARINI A. : Contribuição para o estudo dos Simuliidae (Diptera : Nematocera) da Guiné Portuguesa. Estudo realizado na bacia hidrográfica do Rio Corubal. *Bol. Cult. Guiné Port.*, 1965, 20, 35-65.
- CHARALAMBOUS M., SHELLEY A.J. & GRACIO A.J. dos Santos : Cytogenetical analysis of the *Simulium damnosum* complex (Diptera : Simuliidae) in Guinea Bissau. *Medical and Veterinary Entomology*, in press.
- GRACIO A.J. dos Santos, SHELLEY A.J., RAYBOULD J. & NHAQUE A.T. : The blackflies (Diptera : Simuliidae) of Guinea Bissau. Systematics, distribution and bioecological data. *Acta Parasitológica Portuguesa*, 1994, 2 (1), in press.
- LECUONA M.O. : Nota prévia sobre a existência da oncocercose na Guiné Portuguesa (primeiro caso registado). *An. Inst. Med. trop.*, 1956, 13 (1/2), 83-87.
- LECUONA M.O. : Primeiros dados sobre a distribuição da oncocercose na Guiné Portuguesa. *An. Inst. Med. trop.*, 1959, 16(1/4), 199-208.
- LECUONA M.O. : A oncocercose e o seu interesse médico-social na Guiné Portuguesa. Subsídios para o estudo epidemiológico do foco de Sonaco. *Bol. Cult. Guiné Port.*, 1960, 15 (58), 193-237.
- LOWRY C.A., CHARALAMBOUS M., SHELLEY A.J., GRACIO A.J. dos Santos & HOWARD T.M. : Identification of larvae in the *Simulium damnosum* complex (Diptera : Simuliidae) from Guinea Bissau using multivariate morphometric analysis. In preparation.
- TENDEIRO J. : Estudos sobre simúlios na Guiné Portuguesa. *Garcia de Orta*, 1963, 11 (2), 243-252.

TRANSMISSION OF ONCHOCERCA VOLVULUS, O. OCHENGI AND TYPE D FILARIAE BY SIMULIUM DAMNOSUM S.L. IN NORTH CAMEROON AND ITS INFLUENCE ON THE EPIDEMIOLOGY OF HUMAN ONCHOCERCIASIS

WAHL G.*, ENYONG P.**, SCHIBEL J.M.***, NGOSSO A.****, TUBBESING H.*****, MOYOU R.**, & RENZ A.*****

KEY WORDS : *Onchocerca volvulus*. *Onchocerca ochengi*. *Simulium damnosum*. North Cameroon.

INTRODUCTION

Earlier studies in North-Cameroon showed that the proportion of non- *Onchocerca volvulus* filarial larvae in the onchocerciasis vectors (*Simulium damnosum* s.l.) was

* Institut für Tropenmedizin, Universität Tübingen, Germany.

** Medical Research Station, Kumba, Cameroon.

*** Lehrstuhl für Populationsgenetik, Universität Tübingen, Germany.

**** Hôpital Central, Yaoundé, Cameroon.

***** Hôpital Protestant, Ngaoundéré, Cameroon.

***** Fachgebiet Parasitologie, Universität Hohenheim, Germany.

SITE	SEAS	DAY	DBR	ABR	FLIES DISS (% par)	FLIES L3+ Ov/Oo/TD (% diss fl)	TOTAL L3 Ov/Oo/TD (mL3/inf fl)	DTP Ov/Oo/TD	ATP Ov/Oo/TD
Karna	DS	95	45	16790	6152 (60.4)	34/44/8 (0.6/0.7/0.1)	61/97/8 (1.8/2.2/1.0)	0.5/0.7/0.05	201/128/82
	RS	21	47		1734 (29.5)	10/0/8 (0.6/0/0.5)	23/0/13 (2.3/0/1.6)	0.6/0/0.4	
Galim	DS	17	521	147460	1808 (80.5)	18/124/0 (1.0/6.9/0)	37/552/0 (2.1/4.5/0)	10.9/161.8/0	2446/32722/0
	RS	29	287		3638 (72.5)	14/48/0 (0.4/1.3/0)	31/227/0 (2.2/4.7/0)	2.5/17.5/0	

Table I. – *Simulium damnosum* s.l. biting- and infection rates in a Sudan- (Karna) and Guinea savanna village (Galim) of North-Cameroon.

Ov = *O. volvulus*, Oo = *O. ochengi*, TD = Type D filariae, SEAS = season, DS/RS = dry/rainy season, DAY = number of catching days, FLIES L3+ = flies with L3 in the head, thorax or abdomen, % diss fl = % of dissected flies, mL3/inf fl = mean number of L3 in the head, thorax or abdomen per infected fly, DBR = Daily Biting Rate, ABR = Annual Biting Rate, DTP = Daily Transmission Potential, ATP = Annual Transmission Potential.

Village	Age group	Parasitology					Ophtalmology	
		m/f ex (m/f st)	% snip + (% corr+)	mf/snip	% nod+ (% corr+)	nod/pat	m/f ex (m/f st)	% conch (% corr+)
KARNA	5 - 19 (0-19)	112/125 (110/97)	48.9 %	17.8	10.1 %	1.3	28/13 (46/41)	14.6 %
	20 - 39	27/35 (50/63)	80.6 %	53.9	29.0 %	1.8	26/26 (21/26)	36.5 %
	40 - 59	31/25 (33/39)	82.1 %	46.5	46.4 %	2.5	27/18 (14/16)	68.9 %
	> 59	33/22 (8/7)	90.9 %	82.0	50.9 %	2.6	22/13 (3/3)	82.9 %
	Total	203/207	63.9 % (64.8 %)	56.0	23.4 %	2.1	103/70	49.1 % (32.6 %)
GALIM	5 - 19 (0-19)	44/37 (27/24)	9.9 %	1.7	3.7 %	1.0	10/5 (9/8)	0 %
	20 - 39	1/5 (12/15)	33.3 %	1.8	0 %	–	1/5 (4/5)	0 %
	40 - 59	6/1 (8/10)	14.3 %	5.0	16.7 %	10	5/1 (3/3)	16.7 %
	> 59	4/2 (2/2)	33.3 %	1.5	0 %	–	3/2 (1/1)	0 %
	Total	55/45	13.0 % (17.9 %)	1.9	4.0 %	1.0	19/13	3.1 % (2.9 %)

Table II. – Endemicity of human onchocerciasis in a Sudan- (Karna) and Guinea savanna village (Galim) of North-Cameroon. m/f ex = males/females examined, m/f st = males/females expected in OCP standard population (Moreau *et al.*, 1978), % snip+ = % of examined with one or two positive skin snips, mf/snip = arithmetic mean number of microfilariae per 1 positive snip, % nod+ = % of examined with palpable nodules, nod/pat = a. mean number of nodules per positive patient, oconch = ocular onchocerciasis, % corr+ = percentage corrected for OCP standard population.

astonishingly high (Duke, 1967 ; Renz *et al.*, 1989). Two major forms of larvae were described, Type D larvae (Duke, 1967) and Type G larvae (Renz *et al.*, 1989). While the latter was identified as *O. ochengi* from cattle (Wahl *et al.*, 1991), the former was recently discovered to stem from an *Onchocerca* species in wart hogs (Wahl and Bain, 1993). In an epidemiological field study in two villages in North-Cameroon we assessed the yearly co-transmission of these filariae by man-biting *S. damnosum* s.l. and its possible impact on the local epidemiology of human onchocerciasis.

MATERIAL AND METHODS

The biting and infection rate of *Simulium damnosum* s.l. was assessed during two years near one village in a cattle-raising area in the Guinea Savanna (Galim, estimated population : humans : 350, cattle : 7,000) and another in the Sudan savanna (Karna, 500), where only nomadic cattle (estimated 500) are present from January to April. Flies were caught by volunteers in exhaustors ("sucking tubes") from 7.00 h to 18.00 h at the nearly perennial rivers (Vina du Sud and Benoué) during 1-30 days per month (mean : 6.8, total : 162 days). A proportion of the flies was subsequently dissected, fresh or after deep freezing, in the laboratory. Filarial infective larvae (L3) in the flies were identified morphologically and a proportion of them was sent to Tübingen for identification by specific DNA probes. Sibling species of *S. damnosum* s.l. were identified by cytotaxonomy of larvae in the nearby rivers and morphology of adult flies. The endemicity of human onchocerciasis in the two villages was assessed in random samples of autochthonous volunteers by parasitological examination (2 skin snips from the iliac crest, palpation of nodules) and by thorough ophthalmological examination with a slit lamp. "Ocular onchocerciasis" was defined as sclerosing keratitis, chorioretinitis and/or microfilariae in the cornea and/or anterior chamber.

RESULTS

S. squamosum was the predominant species in the Guinea savanna-site in both rainy and dry season (98.4 % of 63 identified larvae, 97.4 % of 2,970 identified adults). In the Sudan savanna-site the vectors in both seasons were mainly *S. squamosum* and *S. damnosum* s.str. (65.7 % and 32.9 % respectively of 70 larvae, 62.9 % and 37.1 % of 2,606 adults). Both species were attracted 2-3 times more to a cattle than to a human bait. Flies caught on cattle and on man carried all three types of filarial larvae. The infection rates and the proportions of the different filariae were not different in the two vector species and between flies caught on cattle and on man.

The morphological identification of 88 "wild" infective larvae of *O. volvulus*, 160 *O. ochengi* and 29 Type D L3 was confirmed for 90.1 %, 85.7 % and 100 % of the larvae respectively by sequential hybridisation of single L3 with an *O. volvulus*-specific and an *O. ochengi/O. volvulus*-specific DNA probe.

In Karna a moderate Annual Biting Rate (16,800) led to a moderate Annual Transmission Potential of 411 L3 per man per year (Table I). About half of these larvae (48.9 %) were *O. volvulus*, 31.1 % were *O. ochengi* and 20 % Type D. In Galim the biting rate was 8.7 times higher (147,500) and the

ATP 85.6 times higher (35,000) than in Karna. However, only 7 % of all L3 were *O. volvulus*; the remaining ones were *O. ochengi*. Nevertheless, the *O. volvulus*-transmission potential in Galim (2,446) was 12.2 times higher than in Karna (201, Table I).

Onchocerciasis was hyperendemic and severe in Karna with 64.8 % microfilariae carriers, a mean microfilarial load of 56 % per positive snip and 32.6 % of the population showing ocular implications (Table II). In Galim, onchocerciasis was hypoendemic with only 17.9 % skin snip-positives, 1.9 microfilariae per positive snip and 2.9 % ocular onchocerciasis (Table II).

Figure 1. - *Onchocerca* spp. Transmission Potential and Microfilarial Density of *O. volvulus* in a Sudan (Karna) and Guinea savanna village (Galim) of North-Cameroon : Surface areas represent values from Tables I+II and are proportional between villages. Ov = *O. volvulus*, Oo = *O. ochengi*, TD = Type D filariae, ATP = Annual Transmission Potential, Ov-MF/SNIP = arithmetic mean number of *O. volvulus* microfilariae per 1 (positive and negative) snip.

DISCUSSION

The proportion of non-*O. volvulus* filarial larvae in onchocerciasis vectors in North-Cameroon (51.2 - 93 %) is amazingly high. Most of these "animal filariae" (60.9 - 100 %) are *O. ochengi* from cattle. In the rainy season in the Sudan savanna, every third infected *S. damnosum* s.l. carries onchocercal larvae from wart hogs.

This high proportion of animal filariae reflects the high degree of zoophily of the onchocerciasis vectors, which greatly decreases (by the square power) their vector capacity (Garret-Jones, 1964). Furthermore, it must be supposed that many (if not most) of the animal filarial L3 enter man when the vectors bite, and thus provoke an immunological

reaction. In Galim the annual transmission of *O. volvulus* was high (2,500 L3/man) and would normally cause hyperendemic onchocerciasis. However, the mean microfilarial density in the whole population was strikingly low (0.034/snip) and 107 (!) times lower than in Karna (36.29), where the *O. volvulus*-transmission was 12 times lower (Figure 1). This could possibly be due to the very high transmission of *O. ochengi* in Galim (33,000 L3/man x year) which might induce a partial crossreactive immunity in the local population. This hypothesis is strengthened by first immunological studies which demonstrated a high degree of homology between *O. volvulus* and *O. ochengi* in protein profile and serological recognition, and showed significant differences in the serological reactivity between patients from Galim and Karna (Hoch *et al.*, 1992). It is thus concluded that the proximity, throughout the year, of large numbers of cattle infected with *O. ochengi* probably protect the local human population from severe onchocerciasis

ACKNOWLEDGEMENTS

The technical assistance of M. D. Ekale is gratefully acknowledged. This investigation received financial support from the Commission of European Communities (TS2/0184-D (AM) and TS3/CT92-006).

REFERENCES

- DUKE B.O.L. : Infective larvae, other than *Onchocerca volvulus* in *Simulium damnosum*. *Annals of Tropical Medicine and Parasitology*, 1967, 61, 200-205.
- GARRET-JONES C. : The human blood index of malaria vectors in relation to epidemiological assessment. *Bulletin of the World Health Organization*, 1964, 30, 345-362.
- HOCH B., WAHL G., ANYONG P., LÜDER C.G.K., HARNETT W. & RENZ A. : Serological recognition of specific and cross-reactive antigens of *O. ochengi* and *O. volvulus* by infected cattle and humans. *Tropical Medicine and Parasitology*, 1992, 43, 206-207.
- MOREAU J.P., PROST A. & PROD'HON J. : Essai de normalisation de la méthodologie des enquêtes clinico-parasitologiques sur l'onchocercose en Afrique de l'Ouest. *Médecine Tropicale*, 1978, 38, 43-51.
- RENZ A., SCHIBEL J., EICHNER M. & ANYONG P. : Animal filariae in anthrophilic *Simulium* spp. in North Cameroon. *Tropical Medicine and Parasitology*, 1989, 40, 386-387.
- WAHL G., EKALE D., ANYONG P. & RENZ A. : The development of *Onchocerca ochengi* and *O. dukei* microfilariae to infective stage larvae in *Simulium damnosum* s.l. and in members of the *S. medusaeforme* group, following intrathoracic injection. *Annals of Tropical Medicine and Parasitology*, 1991, 85, 324-328.
- WAHL G. & BAIN O. : Development in *Simulium damnosum* s.l. of two *Onchocerca* species from the African wart hog, *O. ramachandri* and *Onchocerca* n. sp. to infective larvae resembling Type D-larvae Duke, 1967. 1993, submitted.

PREPATENCY PERIOD AND SOME ASPECTS OF THE EPIZOOTIOLOGY OF *ONCHOCERCA OCHENGI* INFESTATION IN CATTLE IN THE ADAMAWA PLATEAU, CAMEROON

ACHU-KWT*, DAIBER W.H.** , RENZ A.***, WAHL G.** & WANJI S.****

KEY WORDS : *Onchocerca ochengi*. epizootiology. prepatency. infective larvae. cattle. Cameroon.

SUMMARY

In an onchocerciasis endemic area, calves which were one to 24 months old, were examined for palpable *Onchocerca ochengi* nodules and microfilariae in skin snips. A highly infested bait oxen was used for the production of infective larvae through the vector *Simulium squamosum*. A prepatent period of about 10 months for the appearance of palpable nodules and skin microfilariae was identified, and the prevalence of nodules (80.5 %) and the microfilarial density (0.74 microfilariae per mg skin biopsy) was highest in the 19-24 months old animals. Peaks of 4.8 infective larvae per blood fed fly were reached during infective larvae production. The ease of counting palpable nodules, collecting blood and skin snips, performing nodulectomy, isolating adult worms, keeping these *in vitro* and producing infective larvae provides favorable conditions for the use of this animal model for *in vivo* chemotherapy and vaccine development research.

INTRODUCTION

Although there is a high prevalence of bovine onchocercosis in North Cameroon coupled with a high incidence of multiple-species concurrent infestations (Wahl *et al.*, in press), very little knowledge exists on the biology, immunology and pathology of *Onchocerca ochengi* in particular. There is emphasis on the study of this filaria because its life cycle, infective larvae (Wahl *et al.*, 1991) and nodule formation are very similar to those of *O. volvulus*. The need for expanded research on most of these aspects of bovine onchocercosis has been growing in recent years because of its possible implication in the control (zooprophylaxis and crossreactive concomitant immunity) of human onchocerciasis in a highly endemic region (Renz *et al.*, 1989 ; Wahl, 1991 ; Hoch *et al.*, 1993).

In this survey we investigated the rate of acquisition of palpable nodules and skin microfilariae of *O. ochengi* during natural infestation in cattle with a view to find out the prepatent period of the parasitism. Another objective was to find out how feasible it was to use cattle naturally infested with *O. ochengi* to produce infective larvae which were needed for immunological studies.

* Institute of Animal Research, Wakwa, B.P. 65, Ngaoundere, Cameroon.

** Institut für Tropenmedizin, Wilhelmstrasse 27, 72074 Tübingen, Germany.

*** Fachgebiet Parasitologie, Universität Hohenheim, Emil-Wolff-Strasse 34, 70599 Stuttgart, Germany.

**** Muséum National d'Histoire Naturelle, 61, rue Buffon, 75231 Paris Cédex, France.