PURIFICATION OF A FRACTION OF *GIARDIA LAMBLIA* TROPHOZOITE EXTRACT ASSOCIATED WITH DISACCHARIDASE DEFICIENCIES IN IMMUNE MONGOLIAN GERBILS (*MERIONES UNGUICULATUS*)

MOHAMMED S.R.* & FAUBERT G.M.*

Summary :

The effects on disaccharidase activities of challenging gerbils previously exposed to *Giardia lamblia* with fractions of the crude trophozoite extract were examined. Gel filtration of the soluble extract on a Sephacryl S-200 HR column resulted in 3 fractions : F1, F2 and F3. Only a challenge with fraction F1 (0.1 mg total dose) was found to induce disaccharidase deficiencies. Boiling F1 prior to challenge did not change this effect on the enzyme activities. However, the decreases were not obtained when the total F1 dose was reduced to 0.05 mg. Column chromatography of fraction F1 under dissociating and reducing conditions resulted in 2 further fractions : F1a and F1b. Challenging immune gerbils with F1b led to impairments of disaccharidase activity similar to those obtained with F1. Protein analysis of the crude extract, as well as the fractions of the extract, revealed several high and low molecular weight bands. These findings indicate that a constituent(s) of fraction F1b is the portion of the parasite which induces disaccharidase deficiencies in immune gerbils. This fraction consists of proteins ranging in molecular weight from 32 to 200 kDa. In addition the *G. lamblia* fraction involved in the decreases in enzyme activity is heat-stable.

KEY WORDS : *Giardia lamblia*, disaccharidase, Mongolian gerbils, column chromatography, gel filtration, immunity.

MOTS CLÉS : *Giardia lamblia*, disaccharidase, gerboises mongoles, filtration sur gel, colonne de chromatographie, immunité.

Résumé : PURIFICATION D'UNE FRACTION DE TROPHOZOÏTES DE *GIARDIA LAMBLIA* LIÉE À UNE DIMINUTION DE L'ACTIVITÉ DES DISACCHARIDASES CHEZ LE MÉRION (*MERIONES UNGUICULATUS*) IMMUNISÉ

Les effets de l'inoculation de fractions de trophozoïtes de *Giardia lamblia* sur les activités disaccharidasiques chez les gerboises mongoles précédemment infectées avec *G. lamblia* ont été étudiés. La filtration sur gel Sephacryl S-200 HR a produit 3 fractions : F1, F2 et F3. La fraction F1 est la seule qui ait réduit l'activité des disaccharidases lors d'une dose de rappel (de 0, 1 mg). La fraction F1, amenée à son point d'ébullition, conserve son effet négatif sur l'activité des enzymes. Toutefois, l'activité enzymatique n'est plus affectée par F1 lorsque celle-ci est réduite jusqu'à 0,05 mg. La chromatographie sur colonne de la fraction F1 dans des conditions de dissociation et de réduction a mené à 2 autres fractions : F1a et F1b. Utilisée en dose de rappel chez les gerboises précédemment infectées, la fraction F1b a produit des effets similaires à ceux de F1 sur l'activité des enzymes. L'analyse protéique de l'extrait complet des trophozoïtes et des fractions obtenues par chromatographie révèle plusieurs bandes de poids moléculaires variés. Ces résultats indiquent que la fraction de *Giardia* responsable de la déficience de l'activité des disaccharidases est localisée dans la fraction F1b. Le poids moléculaire des protéines de cette fraction varie de 32 à 200 kDa. De plus, la fraction de *G. lamblia* qui est liée à la diminution des activités disaccharidasiques est stable à la chaleur.

INTRODUCTION

Giardia lamblia is a common cause of diarrhea in day-care centres, institutionalized persons, homosexuals and travellers (Keystone *et al.*, 1978; Brodsky *et al.*, 1974; Schmerin *et al.*, 1978). This flagellated, binucleate parasite adheres to the brush border of the host's enterocytes by means of a ventral sucking disk. Many giardial infections are well tolerated and asymptomatic in human patients. However, when illness is evident, the spectrum of symptoms is wide and ranges from acute, self-limiting gastroenteritis to protracted and debilitating malabsorption (Shandera, 1990).

In giardiasis, it has been established that there can be considerable malfunctioning of the epithelium of the small intestine. For example, fat and vitamin B12 malabsorption has been described in human giardiasis (Hoskins *et al.*, 1967; Wright *et al.*, 1977). Vitamin A deficiency has been found in children infected with *G. lamblia* (Mahalanabis *et al.*, 1979). Also, temporary disaccharidase deficiencies have been well documented in this disease. Buret and co-workers (1990) reported decreases in maltase and sucrase activities in mice, following a primary infection with *G. muris*. It was suggested that these disaccharidase deficiencies were due to a diffuse shortening of brush border microvilli. Studies on *G. lamblia* infections in mice found decreases in several brush border enzymes throughout the course of infection (Nain *et al.*, 1991). The degree of decline in these enzyme activities correlated well with the number of trophozoites in the jejunum, with the most severe reductions in enzyme activity occurring during the peak phase of infection.

* Institute of Parasitology of McGill University, Macdonald Campus, 21,111 Lakeshore Road, Ste. Anne de Bellevue, Québec, Canada H9X 3V9.

Correspondence : Dr. G.M. Faubert, Tel. : (514) 398-7724; Fax : (514) 398-7857. E. mail : GAETAN @ PARASIT.LAN.McGILL.CA.

Using the gerbil animal model, Belosevic and colleagues (1989) observed decreases in disaccharidase activity during both primary and challenge infections with *G. lamblia*. The involvement of the host's immune response in the deficiencies was suggested by the ability of a crude extract of the trophozoites to induce these impairments of enzyme activity in immune gerbils (Belosevic *et al.*, 1989). Recent work in our laboratory has confirmed these previous findings (manuscript in preparation). We also showed that the effect of the extract on disaccharidase activity is dose-dependent.

However, the component(s) of the *G. lamblia* crude extract involved in the disaccharidase deficiencies, as well as the mechanism(s) leading to such reductions in enzyme activity, are unknown. *G. lamblia* is considered to be antigenically complex, yet little is known about the structures and properties of these antigens (Chaudhuri *et al.*, 1988).

Cevallos and Farthing (1992) reported a strain-dependent reduction in disaccharidase activities in *G. lamblia*-infected rats. The antigenic differences between these strains may have led to the variable effects on the enzyme activities. The present study was undertaken in order to purify the *G. lamblia* crude antigen extract and identify the specific fraction responsible for inducing the disaccharidase deficiencies observed in immune gerbils. In addition, we undertook an initial characterization of this fraction. This work shows that a specific fraction of the *G. lamblia* crude extract can be linked to the disaccharidase deficiencies in immune gerbils. This fraction is heat-stable and affects the enzyme activities with a threshold limit.

MATERIALS AND METHODS

PARASITES

G. lamblia trophozoites, WB strain (American Type Culture Collection n° 30957), originally isolated from a symptomatic patient (Smith *et al.*, 1982), were used throughout this study. Trophozoites were cultured axenically in filter-sterilized TYI-S-33 (trypticase, yeast extract, iron, and serum) medium adapted for *G. lamblia* (Gillin and Diamond, 1979), with 10% adult bovine serum (Sigma Chemical Co., St. Louis, U.S.A.), and supplemented with 100 units/ml of penicillin (Sigma) and 100 g/ml of dihydrostreptomycin sulphate salt (Sigma), in the absence of bovine bile. The trophozoites were passaged twice weekly.

ANIMALS

Six to 10 week old male Mongolian gerbils (*Meriones ungukulatus*), originating from Tumblebrook Farms

(West Brookfield, U.S.A.), were used in this study. They were maintained under standard laboratory conditions and provided with laboratory chow and water *ad libitum*. One week after arrival, gerbils were each treated with 15 mg of metronidazole (Rhône-Poulenc, Montreal, Canada), by oral gavage, for three consecutive days and then allowed to rest for 10 days. This treatment was done to ensure that the gerbils were free of intestinal infections. All inoculations were done orally, to unanaesthetized gerbils. In the primary infection, each gerbil was inoculated with 1×10^6 live *G. lamblia* trophozoites.

PREPARATION OF THE CRUDE EXTRACT OF *G. LAMBLIA* TROPHOZOITES

Trophozoites (from a 72-h culture) were dislodged from the walls of culture vessels by immersion in an ice bath for 15-30 min. They were sedimented by centrifugation at 800g for 10 min, at 4 °C, and washed 5 times in sterile phosphate-buffered saline (PBS) (pH 7.2). Trophozoites, suspended in sterile PBS, were then lysed by discontinuous sonication in an ice bath, using a Sonic Dismembrator (Fisher Scientific, Montreal, Canada) at maximum output, for 10 min. The sonicated material was cleared of insoluble matter by centrifuging at 23,300g for 20 min, at 4 °C, and the supernatant collected as the crude soluble extract.

COLUMN CHROMATOGRAPHY

Samples were fractionated by gel filtration chromatography. For this purpose, approximately 80 ml of pre-swollen Sephacryl S-200 HR gel (Pharmacia LKB Biotechnology, Uppsala, Sweden), wet bead diameter 25-75 µm, were packed into a 1.8 × 50 cm glass column (Bio-Rad, Mississauga, Canada). The void volume of the column was determined by applying Blue Dextran 2000 and the column calibrated with known molecular weight (MW) marker of gel filtration proteins (Pharmacia). For each run, at 4 °C, 1-2 ml of sample (with 10% glycerol) was applied to the gel bed and a flow rate of 5 ml/h maintained using a peristaltic pump (Pharmacia). Fractions of 2 ml each were collected and an elution profile obtained by measuring the optical density (O.D.) at 280 nm and plotting these values against the fraction numbers. According to the O.D. values, each peak and trailing eluates were pooled separately. These pooled fractions were then dialyzed, using Spectra/Por 6 Membrane (MW Cut-Off : 1000) (Spectrum Medical Industries Inc., Los Angeles, U.S.A.), against distilled water for 24 h, at 4 °C. Following dialysis, fractions were lyophilized and resuspended in PBS at a concentration of ~0.3 mg/ml. The fractions were stored at -70 °C until used.

SODIUM DODECYL SULFATE-POLYACRYLAMIDE GEL ELECTROPHORESIS (SDS-PAGE)

Samples were electrophoresed in a Mini-PROTEAN II vertical slab cell (Bio-Rad) using the discontinuous system of Laemmli (1970). Protein samples were separated through a 4% stacking gel and 10% separating gel. Bio-Rad molecular weight standards were run simultaneously. The gel was stained with 0.25% Coomassie Brilliant Blue R-250 dye (Bio-Rad).

MEASUREMENT OF INTESTINAL DISACCHARIDASES

Preparation of intestinal homogenate

Homogenates were prepared as previously described by Belosevic and colleagues (1989), with modifications. Briefly, the gerbils were killed and the small intestine removed and divided into three sections. The segments were placed in ice cold distilled water and each was flushed with 50 ml of distilled water to clear it of intestinal debris. The segments were then slit longitudinally and the mucosa scraped off with a glass microscope slide. The mucosa from all three sections were combined, weighed (wet weight), and placed in four volumes of ice cold distilled water. The mucosal scrapings were then homogenized using a Con Torque power unit at maximum speed (Eberbach Corp., Ann Arbor, U.S.A.). Homogenates were stored, without prior centrifugation, at -70 °C until used in the assay.

Assay for intestinal disaccharidases

Disaccharidase activity of homogenated mucosa was measured using the glucose oxidase peroxidase assay of Dahlqvist (1968), as modified by Belosevic *et al.* (1989), which is based on a colour reaction with the glucose liberated by a specific disaccharidase in 60 min. The assay consisted of adding to each well of a 96-well Nunc microwell plate (Gibco BRL, Burlington, Canada) 0.01 ml of appropriately diluted mucosal sample (homogenate) and 0.01 ml substrate-buffer solution of a disaccharidase to be measured, in quadruplicate. The plate was then incubated at 37°C in humidified atmosphere for 60 min. After incubation, 0.3 ml tris-glucose oxidase reagent was added to each well and the plate incubated for an additional 60 min. The plate was then read at 415 nm using a Bio-Tek microplate reader (Mandel Scientific, Guelph, Canada). For each assay, eight wells of reagent blank and a glucose standard series (2, 6, and 10 µg glucose) in quadruplicate were also done. Disaccharidase activity is expressed as units/mg protein in the mucosal sample (U/mg), where units represent µmoles of disaccharide hydrolyzed/min.

Reagents

All chemicals were obtained commercially. D-(+)-

Glucose (G-8270), β-lactose (L-3750), maltose (M-5885), D-(+)-trehalose (T-5251), maleic acid (M-0375), glucose oxidase (G-1262), o-dianisidine (D-9143), peroxidase (P-8000), and triton X-100 (T-6878) were purchased from Sigma. Sucrose (S5-3) was obtained from Fisher Scientific and tris (ultra pure - 819623) was purchased from ICN Biomedicals Canada Ltd. (St. Laurent, Canada).

MEASUREMENT OF PROTEIN CONCENTRATION

Protein concentrations were determined as described by Lowry *et al.* (1951), using 0.2 ml of sample and a final reaction volume of 1.3 ml. For each protein assay, a standard curve was prepared using freshly dissolved bovine serum albumin (BSA).

STATISTICAL ANALYSIS

Statistical significance was determined using the Mann-Whitney *U*-test. Significance was assigned at the probability level of $P < 0.05$.

RESULTS

PURIFICATION OF THE CRUDE EXTRACT OF *G. LAMBLIA* TROPHOZOITES

The crude soluble extract of *G. lamblia* trophozoites was subjected to column chromatography using Sephacryl S-200 HR gel filtration in order to fractionate its components. Three light-absorbing (280 nm) peaks were observed, one of which appeared in the void volume, one in the middle as a shoulder of the first peak, and another at the end of total column volume (Fig. 1). Eluted materials were collected to give fractions F1 ($> 15 \times 10^4$ MW), F2 ($\sim 6.7 \times 10^4$ MW), and F3 ($< 1.4 \times 10^4$ MW), as indicated in Fig. 1. These were pooled from fractions 16-22, 23-29 and 34-43, respectively.

CHALLENGE WITH THE FRACTIONS OF *G. LAMBLIA* SOLUBLE EXTRACT

To determine whether any one of the fractions of the soluble extract (described above) could affect disaccharidase activity in gerbils previously exposed to *G. lamblia*, each animal was challenged with 0.1 mg of F1, F2, or F3. The gerbils were challenged on day 50 post-infection (p.i.), which is well past the elimination of the parasite (around day 30 p.i.), as determined by trophozoite counts from intestinal washings and fecal cyst release. The animals were not treated for the primary infection prior to challenging, since the age-matched control gerbils had also received the

Fig. 1. – Chromatograph of the soluble extract of sonicated *G. lamblia* trophozoites.

Approximately 1.8 mg of the soluble extract were run through a Sephacryl S-200 HR column at a flow rate of 5 ml/h, using a buffer of 0.15 M PBS, 0.02% sodium azide (pH 7.2). Fractions were collected and pooled to give F1, F2 and F3, as indicated. The column was calibrated with Blue Dextran 2000 (BD; $> 200 \times 10^4$ MW), Aldolase (Ald; 15.8×10^4 MW), Bovine Serum Albumin (BSA; 6.7×10^4 MW), and Ribonuclease A (RnA; 1.4×10^4 MW).

Fig. 2. – Disaccharidase activities in gerbils challenged with different fractions of the soluble extract of *G. lamblia* trophozoites. Each gerbil was challenged with 0.1 mg of F1 (▧), F2 (▨) or F3 (▩) 50 days after a primary infection with 1×10^6 live trophozoites. Control gerbils (■) were age-matched and received the primary infection, but were not challenged. Results are expressed as mean $\pm$ s.e., $n = 4$ for unchallenged controls and $n = 5$ for fraction-challenged groups.

* Differences between control and challenged groups were significant, $P < 0.05$.

Fig. 3. – Disaccharidase activities in gerbils challenged with varying amounts of fraction F1. Gerbils were challenged 50 days following a primary infection with 1×10^6 live trophozoites. Control gerbils were age-matched and received a primary infection, but were not challenged. Results are for day 4 post-challenge and are expressed as mean $\pm$ s.e., $n = 4$ for unchallenged controls (■) and $n = 5$ for F1-challenged groups (▨).

* Differences between control and challenged groups were significant, $P < 0.05$.

primary inoculation. The activities of the enzymes lactase, maltase, sucrase, and trehalase were then measured on days 2 and 4 post-challenge (p.c.), as these were the days on which disaccharidase activity was affected during challenge inoculations with the whole crude extract (manuscript in preparation). As shown in Fig. 2, the activity of sucrase was not affected by any of the three fractions of the extract, as levels on both days 2 and 4 p.c. remained comparable to controls. Although the mean activity on day 2 p.c. in F1-challenged animals (0.078 ± 0.018 U/mg) seemed to be lower than in the control (0.12 ± 0.0063 U/mg), this difference was not significant as determined through statistical analysis. However, the activities of lactase, maltase, and trehalase were reduced by as much as 47% on day 4 p.c. in F1-challenged gerbils. In addition, an F1 challenge also led to a decrease in maltase activity on day 2 p.c. On the other hand, challenging immune gerbils with either fraction F2 or F3 had no effect on any of the disaccharidase activities.

In order to determine whether there was a threshold effect of fraction F1 on enzyme activities, immune gerbils were challenged with 0.075 mg or 0.05 mg total of F1. Disaccharidase activity was then measured on day 4 p.c., since this was the time when the most extensive effects on the enzymes were observed with the 0.1 mg F1 challenge. Sucrase activity remained unaffected by this fraction of the soluble extract, regardless of the dosage (Fig. 3). However, lactase

and maltase activities were significantly reduced with an F1 dose of 0.075 mg, compared to unchallenged controls. These decreases in disaccharidase activity were similar to the reduced levels observed in the gerbils challenged with 0.1 mg of F1. There were no reductions in the activities of any of the four enzymes when a dose of 0.05 mg was given to the animals.

In order to determine whether the portion of F1 responsible for inducing the disaccharidase deficiencies is heat-stable, it was boiled at 100°C for 5 min and used to challenge infected gerbils. Decreases in lactase, maltase, and trehalase activities were detected on day 4 p.c., while sucrase activity did not significantly differ from control levels (data not shown). These findings were similar to those obtained with F1 which was not boiled and indicate that the *G. lamblia* fraction involved in the disaccharidase deficiencies is heat-stable.

ISOLATION OF F1 SUB-FRACTIONS

In order to further purify the fraction of the *G. lamblia* extract involved in the disaccharidase deficiencies, fraction F1 was subjected to gel filtration chromatography using a Sephacryl S-200 HR column. The dissociating agent guanidine HCl (Sigma) and the reducing agent DL-dithiothreitol (Sigma) were used as an eluent to disrupt non-covalent interactions and break disulfide bonds in fraction F1. Two peaks were observed, one of which eluted with the void volume

Fig. 4. – Chromatograph of fraction F1. Approximately 2 mg of F1 were applied on a Sephacryl S-200 HR column (1.8×50 cm) at a flow rate of 5 ml/h. The eluent consisted of 2 M guanidine-HCl, 2.5 mM dithiothreitol and 0.02% sodium azide, in PBS (pH 7.2). Fractions were collected and pooled to give F1a and F1b, as indicated. The column was calibrated with Blue Dextran 2000 (BD; $> 200 \times 10^4$ MW), Bovine Serum Albumin (BSA; 6.7×10^4 MW), and Ribonuclease A (RnA; 1.4×10^4 MW).

Fig. 5. – Disaccharidase activities in gerbils challenged with fractions F1a and F1b. Gerbils were each challenged with 0.1 mg of F1a or F1b 50 days following a primary infection with 1×10^6 live trophozoites. Control gerbils were age-matched and received a primary infection, but were not challenged. Results are for as mean $\pm$ s.e., $n = 4$ for unchallenged controls (■) and $n = 5$ for F1a/F1b challenged groups (▨). * Differences between control and challenged groups were significant, $P < 0.05$.

Fig. 6. – SDS-PAGE of crude extract of *G. lamblia* trophozoites and partially purified fractions of the extract. The gel was stained with Coomassie brilliant blue R-250. Molecular weight standards are shown on the left. Lane 1 : crude extract; 2 : F1; 3 : F2; 4 : F3; 5 : F1a; 6 : F1b.

($\sim 200 \times 10^4$ MW) and the other appeared in the high molecular weight ($> 6.7 \times 10^4$ MW) region of the separation (Fig. 4). Fractions 15-19 and 20-25 were collected separately to give pooled fractions F1a and F1b, respectively.

CHALLENGE WITH FRACTIONS F1a AND F1b

To determine whether disaccharidase activity could be affected by one of the fractions of F1, immune gerbils were each challenged with a total of 0.1 mg of either F1a or F1b. Enzyme activities were then measured on day 4 p.c. The activities of all four enzymes remained comparable to control levels in F1a-challenged animals (Fig. 5). However, challenging with fraction F1b resulted in significant decreases in the activities of lactase, maltase, and trehalase. These reductions ranged from 39% to 71%. Although sucrase activity seemed to be lowered by an F1b challenge (0.071 ± 0.014 U/mg compared to a control level of 0.1 ± 0.0075 U/mg), this difference was not significant given the sample sizes.

SDS-PAGE

In order to determine whether there are proteins unique to the fractions affecting enzyme activity, the protein profiles of the crude extract and of the partially purified extract fractions were compared follo-

wing SDS-PAGE. Several high and low molecular weight bands were seen (Fig. 6). The crude extract (lane 1) showed approximately 20 discernable protein bands, with molecular weights ranging from 14 to 130 kDa. However, the banding pattern of fraction F1 (lane 2) was in the 25 to 200 kDa range and F2 (lane 3) was in the smaller region of 30 to 55 kDa. Fraction F3 (lane 4) showed only one protein band, of 20 kDa. For both F1a (lane 5) and F1b (lane 6), a number of bands were detected, mainly in the molecular weight region of 32 to 200 kDa. However, there were two discrete bands seen in fraction F1a which were absent from F1b. These had molecular masses of 25 kDa and 27 kDa (arrows, lane 5). In addition, major bands of 30 kDa and 37 kDa were seen in all samples, with the exception of fraction F3. Both fractions which were implicated in the disaccharidase deficiencies – F1 and F1b – showed several proteins in common with the other fractions.

DISCUSSION

G. lamblia is the most common intestinal protozoan parasite of humans and the acute to chronic diarrhea which it can cause is often associated with intestinal malabsorption (Farthing, 1992). Even when *Giardia* is not suspected, cases of

malabsorption may be due to latent giardiasis, as the organism can be found in the intestine without appearing in feces (Kamath and Murugasu, 1974). However, it has been found that the disaccharidase deficiencies which contribute to the malabsorption of nutrients in giardiasis can occur in infected animal models even in the absence of the live parasite (Belosevic *et al.*, 1989).

In this study, we examined the contribution of the parasite to host malabsorption of disaccharides in giardiasis. Our chromatographic profile of *Giardia* proteins is supported by work done by Chaudhuri and colleagues (1988), who obtained a similar three-peak separation of the *G. lamblia* soluble extract, using Sephacryl S-300 gel filtration. Moreover, these researchers found that maximum antigenic activity was associated with their high molecular weight fraction, which eluted in the void volume and should correspond to fraction F1 in this study. Following separation of the *G. lamblia* soluble extract through Sephacryl S-200 HR chromatography, we were able to identify F1 as the fraction which causes the disaccharidase deficiencies in immune gerbils. The patterns of lactase, maltase and trehalase reductions on days 2 and 4 p.c. in gerbils challenged with F1 mimic those obtained in previous studies with live- and extract-challenged animals (manuscript in preparation). The inability to detect significant differences between control and test groups when measuring sucrase activity may simply be due to the sample sizes. Using a greater number of animals in control and test groups may allow for any differences to be clearly shown, in a statistically significant manner. In addition, there was a threshold effect of fraction F1 on disaccharidase activity and the constituent of F1 involved in the impairments of enzyme activity was found to be heat-stable.

We were able to separate F1 into two fractions, F1a and F1b, again using a Sephacryl S-200 HR column. The use of an eluent containing the dissociating agent guanidine HCl and the reducing agent dithiothreitol, to disrupt non-covalent interactions and break disulfide bonds in F1, allowed for the further separation of giardial proteins. The ability of fraction F1b to induce similar disaccharidase deficiencies in immune gerbils as when challenging with F1 or the crude trophozoite extract (manuscript in preparation), indicates that we have isolated a fraction containing the parasite-specific factor that leads to disaccharide malabsorption. Both inter- and intra-strain antigenic variation among *G. lamblia* isolates have been reported (Smith *et al.*, 1982; Aggarwal and Nash, 1988) and they could contribute to the variability of symptoms seen in giardiasis. Perhaps infections with *Giardia* strains which possess the relevant protein(s) present in F1b lead to disaccharidase deficiencies in the host. Conversely, an

infection with a strain which lacks the key portion of fraction F1b would not affect disaccharide absorption. The complexity of the soluble extract of *G. lamblia* was demonstrated by the SDS-PAGE protein analysis. The present study revealed that approximately 20 protein bands, ranging from 14 to 130 kDa, could be detected. These findings are supported by those of other researchers, who have reported between 20 to 28 protein bands with molecular weights ranging from a low of 10 kDa up to 140 kDa (Moore *et al.*, 1982; Smith *et al.*, 1982; Chaudhuri *et al.*, 1988). We also observed some very faint bands in the region between 130 and 200 kDa (which were concentrated and clearly visible in fraction F1). Such poorly discernable high molecular weight bands in the crude extract have been reported by Smith and colleagues (1982), as well. Given the inability of fraction F2 proteins to affect enzyme activity, it is likely that any protein(s) in F1b responsible for the disaccharidase deficiencies is in the 55 to 200 kDa region.

The role of immune mechanisms in producing malabsorption and intestinal symptoms is unclear. Daniels and Belosevic (1992) found that challenging mice with an extract of *G. muris* led to more depressed disaccharidase activity in susceptible animals as compared to resistant mice. These authors suggested that this finding could be due to differences in the immune response to different parasite antigens. So selective immune recognition of the relevant parasite antigen, present in fraction F1b, could ultimately result in enzyme deficiencies. The specific type of immune response to F1b which could cause a decline in disaccharidase activities remains unknown. Based on other studies on intestinal damage, the disaccharidase deficiencies in giardiasis may be linked to a mucosal mast cell response to an F1b antigen. Curtis *et al.* (1990) demonstrated decreased disaccharidase activity in rats following repeated antigen challenge. These enzyme deficiencies were linked to a mucosal mast cell response, as both mast cell proliferation and degranulation were observed. Since the gerbils used in this study were outbred, it is possible that the variation in enzyme activity observed within a test group could be due to varying degrees of a mast cell immune response. It is important to note that there are possible non-immune mechanisms which could also contribute to disaccharidase deficiencies. For example, *Giardia* trophozoites are known to possess proteinase activity (Hare *et al.*, 1989). A particular proteinase could be present in fraction F1b which affects the enzyme levels as observed. It is apparent, though, that whatever the constituents of F1b which are involved in the disaccharidase deficiencies, they are able to resist the protease activity present in the gut, through an unknown mechanism.

The present study has clearly implicated fraction F1b of the *G. lamblia* trophozoite extract in the disaccharidase deficiencies observed in immune gerbils. This fraction acts with a threshold limit on enzyme activity. It seems probable that without the relevant component of F1b, *Giardia* strains cannot cause disaccharidase deficiencies in infected hosts. However, further investigation of the mechanism(s) of interaction between F1b and the mucosal epithelial cells, including the specific portion of fraction F1b involved in the intestinal dysfunction, is necessary.

ACKNOWLEDGEMENTS

Research at the Institute of Parasitology is supported by the Natural Sciences and Engineering Research Council of Canada and the Fonds FCAR pour l'aide à la recherche.

REFERENCES

- AGGARWAL A. & NASH T.E. Antigenic variation of *Giardia lamblia* in vivo. *Infection and Immunity*, 1988, 56, 1420-1423.
- BELOSEVIC M., FAUBERT G.M. & MACLEAN J.D. Disaccharidase activity in the small intestine of gerbils (*Meriones unguiculatus*) during primary and challenge infections with *Giardia lamblia*. *Gut*, 1989, 30, 1213-1219.
- BRODSKY R.E., SPENCER H.C. & SCHULTZ M.G. Giardiasis in American travelers to Soviet Union. *Journal of Infectious Diseases*, 1974, 130, 319-323.
- BURET A., GALL D.G. & OLSON M.E. Effects of murine giardiasis on growth, intestinal morphology, and disaccharidase activity. *Journal of Parasitology*, 1990, 76, 403-409.
- CEVALLOS A.M. & FARTHING M.J.G. Small intestinal functional damage in experimental giardiasis is strain-dependent. *Gastroenterology*, 1991, 102, A602.
- CHAUDHURI P.P., PAL S., PAL S.C. & DAS P. Studies on *Giardia lamblia* trophozoite antigens using sephacryl S-300 column chromatography, polyacrylamide gel electrophoresis and enzyme-linked immunosorbent assay, in : *Advances in Giardia Research*. Wallis P.M. & Hamrond B.R. (eds), University of Calgary Press, Calgary, 1988, 191-194.
- CURTIS G.H., PATRICK M.K., CATTO-SMITH A.G. & GALL D.G. Intestinal anaphylaxis in the rat : effect of chronic antigen exposure. *Gastroenterology*, 1990, 98, 1558-1566.
- DAHLQVIST A. Assay of intestinal disaccharidases. *Analytical Biochemistry*, 1968, 22, 99-107.
- DANIELS C.W. & BELOSEVIC M. Disaccharidase activity in the small intestine of susceptible and resistant mice after primary and challenge infections with *Giardia muris*. *American Journal of Tropical Medicine and Hygiene*, 1992, 46, 382-390.
- FARTHING M.J.G. New perspectives in giardiasis. *Journal of Medical Microbiology*, 1992, 37, 1-2.
- GILLIN F.D. & DIAMOND L.S. Axenically cultivated *Giardia lamblia* : growth, attachment and the role of L-cysteine, in : *Waterborne Transmission of Giardiasis*. Jakubowski W. & Hoff J.C. (eds), U.S. Environmental Protection Agency, Cincinnati, 1979, 270-272.
- HARE D.F., JARROLL E.L. & LINDMARK D.G. *Giardia lamblia* : characterization of proteinase activity in trophozoites. *Experimental Parasitology*, 1989, 68, 168-175.
- HOSKINS L.C., WINAWER S.J., BROITMAN S.A., GOTTLIEB L.S. & ZAMCHEK N. Clinical giardiasis and intestinal malabsorption. *Gastroenterology*, 1967, 53, 265-279.
- KAMATH K.R. & MURUGASU R. A comparative study of four methods for detecting *Giardia lamblia* in children with diarrheal disease and malabsorption. *Gastroenterology*, 1974, 66, 16-21.
- KEYSTONE J.S., KRAJEN S. & WARREN M.R. Person-to-person transmission of *Giardia lamblia* in day-care nurseries. *Canadian Medical Association Journal*, 1978, 119, 241-248.
- LAEMMLI U.K. Cleavage of structural proteins during assembly of the head of bacteriophage T4. *Nature*, 1970, 227, 680-685.
- LOWRY O.H., ROSEBROUGH N.J., FARR A.L. & RANDALL R.J. Protein measurements with folin phenol reagent. *Journal of Biological Chemistry*, 1951, 193, 265-275.
- MAHALANABIS D., SIMPSON T.W., CHAKRABORTY M.L., GANGULI C., BHARRACHARJEE D.K. & MUKHERJEE K.L. Malabsorption of water miscible vitamin A in children with giardiasis and ascariasis. *American Journal of Clinical Nutrition*, 1979, 32, 313-318.
- MOORE G.W., SOGANDARES-BERNAL F., DENNIS M.V., ROOT D.M., BECKWITH D. & VAN VOORHIS D. Characterization of *Giardia lamblia* trophozoite antigens using polyacrylamide gel electrophoresis, high-performance liquid chromatography, and enzyme-labeled immunosorbent assay. *Veterinary Parasitology*, 1982, 10, 229-237.
- NAIN C.K., DUIT P. & VINAYAK V.K. Alterations in enzymatic activities of the intestinal mucosa during the course of *Giardia lamblia* infection in mice. *Annals of Tropical Medicine and Parasitology*, 1991, 85, 515-522.
- SCHMERIN M.J., JONES T.C. & KLEIN H. Giardiasis : association with homosexuality. *Annals of Internal Medicine*, 1978, 88, 801-803.
- SHANDERA W.X. From Leningrad to the day-care center - The ubiquitous *Giardia lamblia*. *Western Journal of Medicine*, 1990, 153, 154-159.
- SMITH P.D., GILLIN F.D., KAUSHAL N.A. & NASH T.E. Antigenic analysis of *Giardia lamblia* from Afghanistan, Puerto Rico, Ecuador, and Oregon. *Infection and Immunity*, 1982, 36, 714-719.
- SMITH P.D., GILLIN F.D., SPIRA W.M. & NASH T.E. Chronic giardiasis : studies on drug sensitivity, toxin production and host immune response. *Gastroenterology*, 1982, 83, 797-803.
- WRIGHT S.G., TOMKINS A.M. & RIDLEY D.S. Giardiasis : clinical and therapeutic aspects. *Gut*, 1977, 18, 343-350.

Accepté le 21 décembre 1994