RECORD OF *PHLEBOTOMUS (TRANSPHLEBOTOMUS) MASCITTII* GRASSI, 1908 AND *PHLEBOTOMUS (LARROUSSIUS) CHADLII* RIOUX, JUMINER & GIBILY, 1966 FEMALE IN ALGERIA

BERDJANE-BROUK Z.*, CHARREL R.N.** , BITAM I.***, HAMRIOUI B.**** & IZRI A.*

Summary:

We report for the first time the presence of *Phlebotomus mascittii* and the female of *Phlebotomus chadlii* in Algeria. These two species were collected during an entomological study conducted in endemic visceral leishmaniasis focus from the north part of the country, Kabylia.

KEY WORDS: *Phlebotomus mascitti*, *Phlebotomus chadlii*, Algeria.

Résumé : SIGNALEMENT DE *PHLEBOTOMUS (TRANSPHLEBOTOMUS) MASCITTII* GRASSI, 1908 ET DE LA FEMÈLE DE *PHLEBOTOMUS (LARROUSSIUS) CHADLII* RIOUX, JUMINER & GIBILY, 1966 EN ALGÉRIE

Les auteurs signalent pour la première fois la présence de *Phlebotomus mascittii* et de la femelle de *Phlebotomus chadlii* en Algérie. Ces deux espèces ont été capturées lors d'une enquête entomologique réalisée dans un foyer de leishmaniose viscérale du nord du pays, la Kabylie.

MOTS-CLÉS : *Phlebotomus mascitti*, *Phlebotomus chadlii*, Algérie.

Twenty-two phlebotomine sand fly species (Diptera: Psychodidae) have been reported in Algeria, 12 belonging to the *Phlebotomus* genus and 10 to the *Sergentomyia* genus (Belazzoug, 1991). Those included in the *Phlebotomus* genus are of medical importance since they comprise recognized or suspected vectors of leishmaniasis and/or Phlebovirus. We report here for the first time (i) the presence of *Phlebotomus mascittii* in Algeria, and (ii) the presence of the female *Phlebotomus chadlii* in the same area.

The entomological investigation was conducted in Larbaa Nath Iraten (4° 12' 05" E, 36° 38' 10" N at 916 m altitude), in a humid bioclimatic zone, in Kabylia area (Izri *et al.*, 2008). Sand flies collection was performed during summer 2009 using CDC miniature light traps.

A total of 883 sand flies (703 males and 180 females) were captured and morphologically identified during 16 night-CDC traps (55 sand flies/night-CDC traps). Ten distinct species were identified: one species belonging to the *Sergentomyia* genus (*S. minuta*) and nine species to the *Phlebotomus* genus including one female of *P. mascittii* and two females of *P. chadlii* (Table I).

P. mascittii was described in Italy (Roma), then in other countries in the north shore of the Mediterranean basin, from Spain to Turkey (Seccombe *et al.*, 1993). In countries of northern Europe, it was reported in Germany and Switzerland (Naucke *et al.*, 2000). However, *P. mascittii* has always been found in low density. In France, *P. mascittii* species was observed in several departments including in the north, such as Alsace (Callot, 1950). In southern regions, it was usually associated with the main recognized vectors of visceral leishmaniasis, *P. ariasi* and *P. perniciosus* (Rioux *et al.*, 1984; Pesson *et al.*, 1985). It was described as an anthropophylic and aggressive species (Pesson *et al.*, 1985). *P. mascittii* was suspected to be a vector of Mediterranean leishmaniasis, because it was frequently collected from human and dog leishmaniasis in endemic foci (Pesson *et al.*, 1985). However, its vector role has

Specie	Male	Female	Total
<i>P. (Larroussius) perniciosus</i>	564	115	679
<i>P. (Larroussius) longicuspis</i>	84	39	123
<i>P. (Larroussius) langeroni</i>	24	0	24
<i>P. (Larroussius) perfiliewi</i>	3	4	7
<i>P. (Larroussius) ariasi</i>	1	2	3
<i>P. (Larroussius) chadlii</i>	1	2	3
<i>P. (Paraphlebotomus) sergenti</i>	3	7	10
<i>P. (Phlebotomus) papatasi</i>	1		1
<i>P. (Transphlebotomus) mascittii</i>		1	1
<i>S. (Sergentomyia) minuta</i>	22	10	32
Total	703	180	883

Table I. – Sand fly species diversity in LNI, Kabylia area during summer 2009).

* Parasitologie-Mycologie, CHU Avicenne, Université Paris 13, Bobigny, France.

** Unité des Virus Émergents (UMR190), Université de la Méditerranée – Institut de Recherche pour le Développement, Marseille, France.

*** Laboratoire d'Entomologie Médicale, Institut Pasteur d'Alger, Algérie.

**** Laboratoire de Parasitologie, Hôpital Mustapha, Alger, Algérie.

Correspondence: Arezki Izri.

Tel.: 33 (0)1 48 95 54 52 – Fax: 33 (0)1 48 95 56 57.

E-mail: arezki.izri@avc.aphp.fr

Fig. 1. – Spermathecae of *Phlebotomus (Transphlebotomus) mascitti* (photonic microscope $\times 200$).

Fig. 2. – Spermathecae of *Phlebotomus (Larrousius) chadlii* (photonic microscope $\times 200$).

not been confirmed so far. Hence, we noticed for the first time the presence of *P. mascittii* female (Fig. 1) in the southern part of Mediterranean. This female was collected from animal shelter localized in house basement.

P. chadlii was described from Northwest Tunisia (El Kef) among male sand fly specimens. However, the female remained unrecognized until 2006 when it was described in specimens trapped in El Kef, Tunisia (Chamkhi *et al.*, 2006). In Algeria, *P. chadlii* is widely spread in humid, sub humid and arid bioclimatic zones (Dedet *et al.*, 1984). For unknown reasons, in Algeria, only male specimens have been reported so far (Rioux *et al.*, 1970; Dedet *et al.*, 1984). In our survey, three specimens of *P. chadlii* were identified, two females (Fig. 2) and one male, all of them cached in animal shelters. The bioclimatic distribution of *P. chadlii* coincides with that of *P. ariasi* (Dedet *et al.*, 1985), the proven vectors of *L. infantum* and of Sand fly Fever *Phleboviruses* (SFV) in the Mediterranean basin (Izri *et al.*, 2008). Using the mitochondrial *cyt b* gene, Franco *et al.* (2010) reported that *P. chadlii* might be a sister group of the European and the Moroccan *P. ariasi* species. However, to date there is no confirmation neither for their vector role, nor for their trophic preferences. The two females collected in this study were not engorged, thus precluding blood meal analysis. We strongly support the idea of further studies (i) to elucidate the relationship between *P. chadlii* and *P. ariasi*, (ii) to identify their trophic preferences, and (iii) to study the relationship host/leishmaniasis parasite.

ACKNOWLEDGEMENTS

Thanks to Doctor R. Benane, veterinarian in Larbaa Nath Irathen (LNI) region, for their helpful and availability to realize this study.

REFERENCES

- BELAZZOUG S. The sand flies of Algeria. *Parassitologia*, 1991, 33 (Suppl.), 85-87.
- CALLOT J. Présence de *Phlebotomus larrouseii* en Alsace. *Annales de Parasitologie Humaine et Comparée*, 1950, 25 (1-2), 112.
- CHAMKHI J., GUERBOUJ S., BEN ISMAIL R. & GUIZANI I. Description de la femelle de *Phlebotomus (Larroussius) chadlii* Rioux, Juminer et Gibily, 1966 (Diptera : Psychodidae) d'après un exemplaire capturé aux environs du Kef (Tunisie). *Parasite*, 2006, 13 (4), 299-303.
- DEDET J.P., ADDADI K. & BELAZZOUG S. Les phlébotomes (Diptera, Psychodidae) d'Algérie. *Cahier ORSTOM, série Entomologie Médicale et Parasitologie*, 1984, 22, 99-127.
- FRANCO F.A., MORILLAS-MÁRQUEZ F., BARÓN S.D., MORALES-YUSTE M., GÁLVEZ R., DÍAZ V., PESSON B., ALVES-PIRES C., DEPAQUIT J., MOLINA R., AFONSO M.O., GÁLLEGO M., GUERNAOUI S., BOUNAMOUS A. & MARTÍN-SÁNCHEZ J. Genetic structure of *Phlebotomus (Larroussius) ariasi* populations, the vector of *Leishmania infantum* in the western Mediterranean: epidemiological implications. *International Journal of Parasitology*, 2010, 40 (11), 1335-1346.
- IZRI A., TEMMAM S., MOUREAU G., HAMRIOUI B., DE LAMBALLERIE X. & CHARREL R.N. Sand fly fever Sicilian virus, Algeria. *Emerging Infection Diseases*, 2008, 14 (5), 795-797.
- NAUCKE T.J. & PESSON B. Presence of *Phlebotomus (Transphlebotomus) mascittii* Grassi, 1908 (Diptera: Psychodidae) in Germany. *Parasitology Research*, 2000, 86 (4), 335-336.
- PESSON B., LEGER N., MADULO-LEBLOND G., PETAVY A.F. & CAMBON M. Leishmaniasis in Auvergne (France). *Médecine et Maladies Infectieuses*, 1985, 15 (3), 107-109.
- RIOUX J.A., GUY Y., LE COROLLER Y., CROSET H. & ADDADI K. Présence en Algérie de *Phlebotomus (Larroussius) chadlii* Rioux, Juminer et Gibily, 1966. *Bulletin de la Société de Pathologie Exotique*, 1970, 63, 101-104.
- RIOUX J. A., JARRY D. M., LANOTTE G., MAAZOUN R. & KILICK-KENDRICK R. Ecology of leishmaniasis in Southern France. 18. Enzymatic identification of *Leishmania infantum* Nicolle, 1908, isolated from *Phlebotomus ariasi* Tonnoir, 1921, spontaneously infected in the Cévennes. *Annales de Parasitologie Humaine et Comparée*, 1984, 59 (4), 331-333.
- SECCOMBE A.K., READY P.D. & HUDDLESTON L.M. A Catalogue of Old World Phlebotomine Sand flies (Diptera: Psychodidae, Phlebotominae). Occ. Pap. syst. Ent No. 8, The Natural History Museum – London (ed.), Intercept, Andover, 1993.

Received on May 16th, 2011
Accepted on August 25th, 2011