

## Anthelmintic properties of traditional African and Caribbean medicinal plants: identification of extracts with potent activity against *Ascaris suum* *in vitro*

Andrew R. Williams<sup>1,\*</sup>, Jens Soelberg<sup>2</sup>, and Anna K. Jäger<sup>2</sup>

<sup>1</sup> Department of Veterinary Disease Biology, Faculty of Health and Medical Sciences, University of Copenhagen, 1870 Copenhagen, Denmark

<sup>2</sup> Department of Drug Design and Pharmacology, Faculty of Health and Medical Sciences, University of Copenhagen, 2100 Copenhagen, Denmark

Received 6 April 2016, Accepted 2 June 2016, Published online 14 June 2016

**Abstract** – Ascariasis affects more than 1 billion people worldwide, mainly in developing countries, causing substantial morbidity. Current treatments for *Ascaris* infection are based on mass drug administration (MDA) with synthetic anthelmintic drugs such as albendazole, however continual re-infection and the threat of drug resistance mean that complementary treatment options would be highly valuable. Here, we screened ethanolic extracts from 29 medicinal plants used in Africa (Ghana) and the Caribbean (US Virgin Islands) for *in vitro* anthelmintic properties against *Ascaris suum*, a swine parasite that is very closely related to the human *A. lumbricoides*. A wide variety of activities were seen in the extracts, from negligible to potent. Extracts from *Clausena anisata*, *Zanthoxylum zanthoxyloides* and *Punica granatum* were identified as the most potent with EC<sub>50</sub> values of 74, 97 and 164 µg/mL, respectively. Our results encourage further investigation of their use as complementary treatment options for ascariasis, alongside MDA.

**Key words:** *Ascaris suum*, Anthelmintic, *Clausena anisata*, *Zanthoxylum zanthoxyloides*, *Punica granatum*.

**Résumé** – Propriétés anthelminthiques de plantes médicinales traditionnelles d’Afrique et des Caraïbes: identification des extraits ayant une activité puissante *in vitro* contre *Ascaris suum*. L’ascaridiose affecte plus d’un milliard de personnes dans le monde, principalement dans les pays en développement, causant une importante morbidité. Les traitements actuels contre les infections par *Ascaris* se basent sur l’administration massive de médicaments (AMM) avec des anthelminthiques de synthèse comme l’albendazole, mais les réinfections continues et la menace de résistance aux médicaments signifient que des options de traitements complémentaires seraient très utiles. Dans cet article, nous avons criblé des extraits éthanoliques de vingt-neuf plantes médicinales utilisées en Afrique (Ghana) et aux Caraïbes (Îles Vierges Américaines) pour leurs propriétés anthelminthiques *in vitro* contre *Ascaris suum*, un parasite du porc qui est très proche de l’espèce *A. lumbricoides* de l’homme. Une grande variabilité d’activités a été trouvée parmi les extraits, allant de négligeable à puissante. Les extraits de *Clausena anisata*, *Zanthoxylum zanthoxyloides* et *Punica granatum* ont été identifiés comme les plus puissants avec des valeurs de CE<sub>50</sub> de 74, 97 et 164 µg/mL, respectivement. Nos résultats encouragent à plus de recherche sur leur utilisation comme options de traitement complémentaires de l’ascaridiose, aux côtés de l’AMM.

### Introduction

Soil-transmitted helminths remain one of the largest burdens on global health. Altogether, *Ascaris lumbricoides* (roundworm), *Trichuris trichiura* (whipworm) and *Necator americanus* (hookworm) infect more than a billion people, mainly in the developing world [20, 24]. Of these, the most prevalent is *A. lumbricoides*, which was estimated to infect

around 800 million people in 2010, resulting in more than a million years lived with disability (YLD) [20]. Ascariasis can result in malnutrition and inhibit cognitive development and learning in children, and can also interfere with effective expression of immunity to other pathogens and vaccines due to polarisation of the immune system towards a regulatory/Th2-skewed state [6, 21].

At present, *A. lumbricoides* infections are treated through mass drug administration (MDA) programmes, involving annual or bi-annual treatment of school children, mainly with

\*Corresponding author: [arw@sund.ku.dk](mailto:arw@sund.ku.dk)

albendazole [12]. However, the sustainability of this approach has been questioned due to continual re-infection, arising from the hard-shelled eggs which survive for many years in the environment, and the ever-present threat of drug resistance, where cautionary tales can be drawn from the critical levels of resistance that have arisen in helminths of veterinary importance due to mass administration of synthetic drugs [11, 13, 23].

The use of indigenous medicinal plants for control of internal parasites has been practised for centuries, however scientific validation of these traditional practices has been lacking [8, 22]. Thus, a potentially vast resource of natural plant compounds with anthelmintic activity has been underutilised. The use of natural plant extracts, which can be prepared by decoction or other simple procedures, has several advantages such as low cost and easy integration into local community practices, if and where plants are locally available. The rational use of such an approach may complement current MDA programmes and be a useful tool to slow the onset of drug resistance.

Screening for anthelmintic activity is often accomplished by using the free-living nematode *Caenorhabditis elegans* [5, 9] due to low cost and amenability of laboratory culture methods. However, many differences exist between free-living nematodes and parasites such as *Ascaris*, as well as between free-living and parasitic stages of the same species [17]. Thus, anthelmintic activity is ideally assessed using as close as possible a model to the eventual target. *Ascaris suum* is a swine parasite and closely related parasite to *A. lumbricoides*, being morphologically indistinguishable, and indeed, for many years the human and pig forms of *Ascaris* were considered to be a single species [16]. This means that *A. suum* is an excellent model for studying possible new interventions such as vaccines or new drug candidates against *A. lumbricoides*. In particular, *in vitro* studies with *A. suum* are easily performed due to the ability to generate large numbers of infective, parasitic third-stage larvae (L3) from embryonated eggs recovered from the uteri of adult female worms. We have previously used a combination of motility and migration inhibition assays to test for *in vitro* anthelmintic effects of a number of compounds against *A. suum* L3 [25]. Here, we utilised local ethno-medical knowledge to compile a library of traditional medicinal plants from Ghana and the Caribbean, and used *A. suum* L3 to screen > 30 extracts for *in vitro* activity. Our results indicate that some of these plants may have value as a complementary treatment option for ascariasis.

## Materials and methods

### Plant material and extraction

Medicinal plants were collected during the period November 2013 to January 2014 in the Greater Accra region of Ghana or October to November 2014 on the US Virgin Islands (Table 1). The plants were dried at ambient temperature in the shade. The plants were identified and authenticated by an ethnobotanist (Jens Soelberg, University of Copenhagen, Denmark). Botanical voucher specimens were deposited at the Herbaria at the University of Copenhagen, Denmark; and at the University of Ghana, Ghana, and St. George Estate

Botanical Garden, respectively. Voucher numbers are given in Table 1. Ethanolic extracts were prepared by extracting 2 g of ground plant material with 20 mL of 96% ethanol in an ultrasonic bath for 30 min. Extracts were filtered through filter paper and taken to dryness at room temperature under nitrogen. For anthelmintic assays, extracts were dissolved in 100% DMSO.

### Parasite material

Gravid *Ascaris suum* worms were collected from fresh pig intestines at a local slaughterhouse (Danish Crown, Ringsted, Denmark). Eggs were isolated from the uteri of the worms and then subsequently embryonated at 25 °C for at least 60 days in 0.1 M H<sub>2</sub>SO<sub>4</sub>. Full embryonation was confirmed by light microscopy. To obtain the L3, eggs were washed, suspended in Hanks' Buffered Salt Solution (HBSS) and then hatched by stirring together with 2 mm glass beads for 30 min at 37 °C. Viable L3 were then separated from unhatched eggs and debris by overnight migration into sterile HBSS using a Baermann apparatus equipped with 20 µm mesh. The L3 were then washed, counted and suspended in larval culture media (RPMI 1640 supplemented with 2 mM L-glutamine, 100 U/mL penicillin and 100 µg/mL streptomycin) for use in the migration inhibition assay.

### Larval migration inhibition assay

The migration inhibition assay was conducted essentially as described previously [25]. Briefly, 100 L3 (in triplicate) were added to wells on a tissue culture plate and plant extracts in DMSO added (DMSO concentration never exceeded 1%). All assays included 1% DMSO in culture media as a negative control and 50 µg/mL ivermectin (Sigma-Aldrich) as a positive control. The plates were then incubated overnight at 37 °C in 5% CO<sub>2</sub> in air. Then, an equal amount of 1.6% agar solution (45 °C) was added to each well and mixed thoroughly. The agar was allowed to solidify, before fresh culture media were added on top to cover the agar and the plates returned to the incubator overnight. The next day, the media were collected from each well and the number of larvae that had migrated from the setting agar was enumerated by light microscopy.

### Data analysis and statistics

Migration inhibition was calculated relative to larvae incubated only in culture media (+1% DMSO) and expressed as percentage inhibition. Half-maximal effective concentration (EC<sub>50</sub>) values were calculated using non-linear regression. Analyses were performed in GraphPad Prism (v6.00, GraphPad Software, La Jolla, California, USA, [www.graphpad.com](http://www.graphpad.com)).

## Results and discussion

Screening of the plant extracts at a concentration of 1 mg/mL revealed a wide variety of potencies (Fig. 1).

**Table 1.** Names and characteristics of plant species tested for anthelmintic activity against *Ascaris suum*. Plants were collected from either Ghana (G) or the US Virgin Islands (VI).


Species name	Location	Family	Voucher	Plant part
<i>Aloe vera</i> (L.) Burm. f.	VI	Xanthorrhoeaceae	JS 617	Leaves
<i>Boerhavia diffusa</i> L.	G	Nyctaginaceae	JS 281	Aerial parts
<i>Boerhavia erecta</i> L.	G	Nyctaginaceae	JS 282	Aerial parts
<i>Clausena anisata</i> (Willd.) Hook. f. ex Benth.	G	Rutaceae	JS 214	Roots
<i>Deinbollia pinnata</i> Schum. & Thonn.	G	Sapindaceae	JS 202	Aerial parts
<i>Erythrina senegalensis</i> DC.	G	Leguminosae	JS 231	Bark
<i>Flacourtia flavescens</i> Willd.	G	Salicaceae	JS 249	Leaves
<i>Flueggea virosa</i> (Roxb. ex Willd.) Royle	G	Phyllanthaceae	JS 252	Leaves
<i>Gardenia ternifolia</i> Schumach. & Thonn.	G	Rubiaceae	JS 246	Leaves
<i>Gymnanthemum coloratum</i> (Willd.) H. Rob. & B. Kahn	G	Asteraceae	JS 268	Leaves/flowers
<i>Gymnanthemum coloratum</i> (Willd.) H. Rob. & B. Kahn	G	Compositae	JS 268	Roots
<i>Launaea taraxacifolia</i> (Willd.) Amin ex C. Jeffrey	G	Compositae	JS 212	Leaves
<i>Mallotus oppositifolius</i> (Geiseler) Müll. Arg.	G	Euphorbiaceae	JS 208	Leaves
<i>Mucuna pruriens</i> (L.) DC.	VI	Leguminosae	JS 656	Fruit
<i>Newbouldia laevis</i> (P. Beauv.) Seem.	G	Bignoniaceae	JS 216	Leaves
<i>Opuntia</i> sp.	VI	Cactaceae	JS 672	Stem
<i>Paullinia africana</i> R.Br. ex Tedlie	G	Sapindaceae	JS 219	Aerial parts
<i>Phyllanthus amarus</i> Schumach. & Thonn.	G	Phyllanthaceae	JS 237	Aerial parts
<i>Premna quadrifolia</i> Schumach. & Thonn.	G	Lamiaceae	JS 283	Aerial parts
<i>Psidium guajava</i> L.	VI	Myrtaceae	JS 623	Leaves
<i>Punica granatum</i> L.	VI	Lythraceae	JS 615	Fruit peel
<i>Pupalia lappacea</i> (L.) Juss.	G	Amaranthaceae	JS 239	Aerial parts
<i>Rivina humilis</i> L.	VI	Phytolaccaceae	JS 608	Aerial parts
<i>Senna occidentalis</i> (L.) Link	G	Leguminosae	JS 234	Aerial parts
<i>Senna occidentalis</i> (L.) Link	G	Leguminosae	JS 234	Roots
<i>Spathodea campanulata</i> P. Beauv.	G	Bignoniaceae	JS 230	Bark
<i>Spigelia anthelmia</i> L.	VI	Loganiaceae	JS 651	Aerial parts
<i>Stylosanthes erecta</i> P. Beauv.	G	Leguminosae	JS 271	Aerial parts
<i>Thonningia sanguinea</i> Vahl	G	Balanophora	JS 296	Aerial parts
<i>Triumfetta semitriloba</i> Jacq.	VI	Malvaceae	JS 658	Aerial parts
<i>Zanthoxylum zanthoxyloides</i> (Lam.) Zepern. & Timler	G	Rutaceae	JS 243	Roots
<i>Zanthoxylum zanthoxyloides</i> (Lam.) Zepern. & Timler	G	Rutaceae	JS 243	Root bark

A number of extracts had no or negligible activity. However, 10 extracts inhibited larval migration by at least 50%. An arbitrary cut-off value of 90% migration inhibition was chosen to define plants that had potent activity, comparable to the inhibition achieved by 50 µg/mL ivermectin (positive control). Using this criterion, four extracts – *Clausena anisata*, *Zanthoxylum zanthoxyloides* (both the roots and root bark) and *Punica granatum* – were selected on the basis of potent activities for dose-dependent studies to confirm their activities and determine EC<sub>50</sub> values. All four extracts displayed dose-dependent activity (Fig. 2). The EC<sub>50</sub> (95% CI) values for *C. anisata*, *Z. zanthoxyloides* roots, *Z. zanthoxyloides* root bark and *P. granatum* were 74 (63.3–86.8), 97 (63.9–149.3), 132 (105.9–164.6) and 164 (124.7–271.1) µg/mL, respectively.


We have thus confirmed that a number of plants that are traditionally used in medicinal form in *Ascaris*-endemic regions have direct anthelmintic activity against *A. suum*. The identification of three extracts with particularly potent activity is consistent with previous reports concerning their ethno-medical usage and anthelmintic properties. *C. anisata* is used as an anthelmintic by traditional healers in Kenya [24], and has also been noted to have *in vitro* activity against

free-living larvae of the sheep nematode *Haemonchus contortus* [23]. Similarly, *in vitro* studies with *Z. zanthoxyloides* have demonstrated anthelmintic effects of crude extracts from this plant against *H. contortus* and another sheep nematode, *Trichostrongylus colubriformis* [4, 10]. Moreover, the distilled essential oils from *Z. zanthoxyloides* have been shown to have *in vitro* activity against the rat helminth *Strongyloides ratti* [18]. *Punica granatum* extracts also have *in vitro* activity against free-living worms such as *Allolobophora caliginosa* [7] and veterinary helminths such as the poultry roundworm *Ascaridia galli* [3]. Our data, which indicate a strong anti-*Ascaris* effect of these three plant extracts, extend on these previous studies by suggesting that they may have potential for treatment of *A. lumbricoides* in humans.

The active compounds of these extracts were not investigated here, but all three extracts (*Z. zanthoxyloides*, *P. granatum* and *C. anisata*) are known to be rich in secondary compounds such as tannins (with *P. granatum* being a particularly rich source of ellagitannins and gallic acid), flavonoids, terpenes and alkaloids, and these have been implicated in the apparent anti-microbial and ethno-medical properties of these plants [1, 14, 19]. Previous *in vitro* studies with *H. contortus*



**Figure 1.** Inhibition of migratory ability of *Ascaris suum* third-stage larvae after exposure to extracts of medicinal plants (1 mg/mL) or ivermectin (50  $\mu$ g/mL). Results are the mean ( $\pm$  SEM) of three replicates from a single experiment. The vertical dashed line indicates 90% inhibition of larval migration.



**Figure 2.** Dose-dependent inhibition of migration of *Ascaris suum* third-stage larvae after exposure to extracts of three different medicinal plants. Results are the mean of three independent experiments, each performed in triplicate, with the results expressed as the mean  $\pm$  inter-experiment SEM.

and *S. ratti* have suggested that the anthelmintic properties of *Z. zanthoxyloides* are partially dependent on flavonoids and related polymeric tannins, but that other bioactive

compounds such as terpenes derived from the essential oil may contribute to the activity [4, 18]. Furthermore, a related plant, *Z. liebmannianum*, has also been shown to have activity against *A. suum* *in vitro* and here  $\alpha$ -sanshool was isolated as a putative active compound [15]. Ellagitannins or related phenolic compounds found in *P. granatum* have been speculated to be responsible for activity that has been observed against protozoan parasites such as *Cryptosporidium parvum* [2]. Fractionation and identification of the active compounds in these extracts is ongoing in our laboratory.

The confirmation of anti-*Ascaris* activity of some of these traditional extracts encourages refinement of their use as treatments for roundworm infection in areas where these plants are found locally. Whilst our current experiments were performed with third-stage larvae, which are present in the intestinal tract for only a short period of time following egg ingestion, we have shown previously that activity of bioactive compounds against L3 correlates very closely with their activity against both fourth-stage larvae [25] and adult worms (A.R. Williams, unpublished data). Thus, there appears to be clear scope for these remedies to be used as therapeutic treatments against established infections in the small intestine. Having now confirmed the activity of these traditional extracts, further studies to investigate the pharmacokinetics in order to determine the best possible dosage and administration methods of these

extracts are now warranted. Moreover, if activity can be solely or mainly ascribed to a single compound, then the possibility remains of synthesis and production of a new anthelmintic drug.

In conclusion, we have demonstrated that extracts of *C. anisata*, *Z. zanthoxyloides* and *P. granatum* have potent anthelmintic activity against *A. suum* *in vitro*, which encourages further investigation of their use as therapeutic agents against *Ascaris* infections in endemic regions.

**Acknowledgements.** The authors are grateful to Katrine Krydefelt for assistance with extraction of plant material. ARW was supported by the Danish Council for Independent Research (Technology and Production Sciences, Grant No. 12-126630).

## References

1. Agyepong N, Agyare C, Adarkwa-Yiadom M, Gbedema SY. 2014. Phytochemical investigation and anti-microbial activity of *Clausena Anisata* (Willd), Hook. African Journal of Traditional, Complementary, and Alternative Medicines, 11(3), 200–209.
2. Al-Mathal EM, Alsalem AA. 2013. Pomegranate (*Punica granatum*) peel is effective in a murine model of experimental *Cryptosporidium parvum* ultrastructural studies of the ileum. Experimental Parasitology, 134(4), 482–494.
3. Ali N, Jamil A, Shah S, Shah I, Ahmed G, Junaid M, Ahmed Z. 2015. Parasitocidal and brine shrimp cytotoxicity potential of crude methanolic extract of rind of *Punica granatum* Linn against round worms and tape worms. Pakistan Journal of Pharmaceutical Sciences, 28(3), 959–962.
4. Azando EVB, Hounzangbé-Adoté MS, Olounladé PA, Brunet S, Fabre N, Valentin A, Hoste H. 2011. Involvement of tannins and flavonoids in the *in vitro* effects of *Newbouldia laevis* and *Zanthoxylum zanthoxyloides* extracts on the exsheathment of third-stage infective larvae of gastrointestinal nematodes. Veterinary Parasitology, 180(3–4), 292–297.
5. Buckingham SD, Partridge FA, Sattelle DB. 2014. Automated, high-throughput, motility analysis in *Caenorhabditis elegans* and parasitic nematodes: applications in the search for new anthelmintics. International Journal for Parasitology: Drugs and Drug Resistance, 4(3), 226–232.
6. Cooper PJ, Chico ME, Sandoval C, Espinel I, Guevara A, Kennedy MW, Urban JF, Griffin GE, Nutman TB. 2000. Human infection with *Ascaris lumbricoides* is associated with a polarized cytokine response. Journal of Infectious Diseases, 182(4), 1207–1213.
7. Dkhil MA. 2013. Anti-coccidial, anthelmintic and antioxidant activities of pomegranate (*Punica granatum*) peel extract. Parasitology Research, 112(7), 2639–2646.
8. Githiori JB, Höglund J, Waller PJ. 2005. Ethnoveterinary plant preparations as livestock dewormers: practices, popular beliefs, pitfalls and prospects for the future. Animal Health Research Reviews, 6(01), 91–103.
9. Holden-Dye L, Walker RJ. 2014. Anthelmintic drugs and nematicides: studies in *Caenorhabditis elegans*, in WormBook, Editor. C. elegans Research Community, WormBook, DOI: 10.1895/wormbook.1.143.2, <http://www.wormbook.org>
10. Hounzangbe-Adote S, Fouraste I, Moutairou K, Hoste H. 2005. *In vitro* effects of four tropical plants on the activity and development of the parasitic nematode, *Trichostrongylus colubriformis*. Journal of Helminthology, 79(01), 29–33.
11. Jia T-W, Melville S, Utzinger J, King CH, Zhou X-N. 2012. Soil-transmitted helminth reinfection after drug treatment: a systematic review and meta-analysis. PLoS Neglected Tropical Diseases, 6(5), e1621.
12. Keiser J, Utzinger J. 2008. Efficacy of current drugs against soil-transmitted helminth infections: systematic review and meta-analysis. Journal of the American Medical Association, 299(16), 1937–1948.
13. Kotze AC, Hunt PW, Skuce P, von Samson-Himmelstjerna G, Martin RJ, Sager H, Krücken J, Hodgkinson J, Lespine A, Jex AR, Gilleard JS, Beech RN, Wolstenholme AJ, Demeler J, Robertson AP, Charvet CL, Neveu C, Kaminsky R, Rufener L, Alberich M, Menez C, Prichard RK. 2014. Recent advances in candidate-gene and whole-genome approaches to the discovery of anthelmintic resistance markers and the description of drug/receptor interactions. International Journal for Parasitology: Drugs and Drug Resistance, 4(3), 164–184.
14. Mena P, Ascacio-Valdés JA, Gironés-Vilaplana A, Del Rio D, Moreno DA, García-Viguera C. 2014. Assessment of pomegranate wine lees as a valuable source for the recovery of (poly)phenolic compounds. Food Chemistry, 145, 327–334.
15. Navarrete A, Hong E. 1996. Anthelmintic properties of  $\alpha$ -sanshool from *Zanthoxylum liebmannianum*. Planta Medica, 62(03), 250–251.
16. Nejsum P, Betson M, Bendall RP, Thamsborg SM, Stothard JR. 2012. Assessing the zoonotic potential of *Ascaris suum* and *Trichuris suis*: looking to the future from an analysis of the past. Journal of Helminthology, 86(02), 148–155.
17. O'Grady J, Kotze AC. 2004. *Haemonchus contortus*: *in vitro* drug screening assays with the adult life stage. Experimental Parasitology, 106(3–4), 164–172.
18. Olounladé PA, Azando EVB, Hounzangbé-Adoté MS, Ha TBT, Leroy E, Moulis C, Fabre N, Magnaval JF, Hoste H, Valentin A. 2011. *In vitro* anthelmintic activity of the essential oils of *Zanthoxylum zanthoxyloides* and *Newbouldia laevis* against *Strongyloides ratti*. Parasitology Research, 110(4), 1427–1433.
19. Patino L, Prieto RJA, Cuca SLE. 2012. *Zanthoxylum* Genus as a potential source of bioactive compounds, in Bioactive compounds in phytomedicine. Rasooli I, EditorIntech: Rijeka, Croatia. p. 185–218.
20. Pullan R, Smith J, Jasrasaria R, Brooker S. 2014. Global numbers of infection and disease burden of soil transmitted helminth infections in 2010. Parasites & Vectors, 7(1), 37.
21. Steenhard NR, Jungersen G, Kokotovic B, Beshah E, Dawson HD, Urban JF Jr, Roepstorff A, Thamsborg SM. 2009. *Ascaris suum* infection negatively affects the response to a *Mycoplasma hyopneumoniae* vaccination and subsequent challenge infection in pigs. Vaccine, 27(37), 5161–5169.
22. Tolossa K, Debela E, Athanasiadou S, Tolera A, Ganga G, Houdijk J. 2013. Ethno-medicinal study of plants used for treatment of human and livestock ailments by traditional healers in South Omo, Southern Ethiopia. Journal of Ethnobiology and Ethnomedicine, 9(1), 32.
23. Vercruyse J, Albonico M, Behnke JM, Kotze AC, Prichard RK, McCarthy JS, Montresor A, Levecke B. 2011. Is anthelmintic resistance a concern for the control of human soil-transmitted helminths? International Journal for Parasitology: Drugs and Drug Resistance, 1(1), 14–27.

24. WHO. 2012. Accelerating work to overcome the global impact of neglected tropical diseases – a roadmap for implementation. World Health Organization: Geneva.
25. Williams AR, Fryganas C, Ramsay A, Mueller-Harvey I, Thamsborg SM. 2014. Direct anthelmintic effects of condensed tannins from diverse plant sources against *Ascaris suum*. PLoS One, 9(5), e97053.

**Cite this article as:** Williams AR, Soelberg J & Jäger AK: Anthelmintic properties of traditional African and Caribbean medicinal plants: identification of extracts with potent activity against *Ascaris suum* *in vitro*. Parasite, 2016, 23, 24.



An international open-access, peer-reviewed, online journal publishing high quality papers on all aspects of human and animal parasitology

Reviews, articles and short notes may be submitted. Fields include, but are not limited to: general, medical and veterinary parasitology; morphology, including ultrastructure; parasite systematics, including entomology, acarology, helminthology and protistology, and molecular analyses; molecular biology and biochemistry; immunology of parasitic diseases; host-parasite relationships; ecology and life history of parasites; epidemiology; therapeutics; new diagnostic tools.

All papers in Parasite are published in English. Manuscripts should have a broad interest and must not have been published or submitted elsewhere. No limit is imposed on the length of manuscripts.

**Parasite** (open-access) continues **Parasite** (print and online editions, 1994-2012) and **Annales de Parasitologie Humaine et Comparée** (1923-1993) and is the official journal of the Société Française de Parasitologie.

Editor-in-Chief:  
Jean-Lou Justine, Paris

Submit your manuscript at  
<http://parasite.edmgr.com/>